



Endogenous and Borrowed Proteolytic Activity in the *Borrelia*

James L. Coleman,^{a,c} Jorge L. Benach,^{a,c}  A. Wali Karzai^{a,b,c}

^aDepartment of Microbiology and Immunology, Stony Brook University, Stony Brook, New York, USA

^bDepartment of Biochemistry and Cell Biology, Stony Brook University, Stony Brook, New York, USA

^cCenter for Infectious Diseases, Stony Brook University, Stony Brook, New York, USA

SUMMARY	1
INTRODUCTION	2
Serine Proteases	2
Threonine Proteases	3
Cysteine Proteases	3
Aspartyl Proteases	3
Metalloproteases	3
ENDOGENOUS PROTEASES OF THE BORRELIA	3
Proteolytic Component of <i>Borrelia</i>	3
Amino and Carboxypeptidases of <i>B. burgdorferi</i>	5
FtsH of <i>B. burgdorferi</i>	6
Lon Proteases of <i>B. burgdorferi</i> Are Unusual in the Eubacteria	7
HtrA of <i>Borrelia</i> , a New Strategy for Pathogenesis	8
BORRELIA AND BORROWED PROTEOLYSIS	11
Plasminogen Activation System	11
Plasmin(ogen) Binding and Invasion by Bacteria	12
Plasmin(ogen) Binding by <i>Borrelia</i>	12
Substrates Degraded by <i>Borrelia</i> with Bound Plasmin	13
<i>Borrelia</i> and Plasminogen Receptors	13
Interaction of <i>Borrelia</i> with Other PAS Components	15
Borrowed Proteolysis and Host Defense	16
CONCLUDING REMARKS	16
ACKNOWLEDGMENTS	17
REFERENCES	18

SUMMARY The *Borrelia* spp. are tick-borne pathogenic spirochetes that include the agents of Lyme disease and relapsing fever. As part of their life cycle, the spirochetes traffic between the tick vector and the vertebrate host, which requires significant physiological changes and remodeling of their outer membranes and proteome. This crucial proteome resculpting is carried out by a diverse set of proteases, adaptor proteins, and related chaperones. Despite its small genome, *Borrelia burgdorferi* has dedicated a large percentage of its genome to proteolysis, including a full complement of ATP-dependent proteases. Energy-driven proteolysis appears to be an important physiological feature of this dual-life-cycle bacterium. The proteolytic arsenal of *Borrelia* is strategically deployed for disposal of proteins no longer required as they move from one stage to another or are transferred from one host to another. Likewise, the *Borrelia* spp. are systemic organisms that need to break down and move through host tissues and barriers, and so their unique proteolytic resources, both endogenous and borrowed, make movement more feasible. Both the Lyme disease and relapsing fever *Borrelia* spp. bind plasminogen as well as numerous components of the mammalian plasminogen-activating system. This recruitment capacity endows the spirochetes with a borrowed proteolytic competency that can lead to increased invasiveness.

KEYWORDS *Borrelia*, borrowed proteolysis, plasminogen, proteases, proteolytic enzymes

Citation Coleman JL, Benach JL, Karzai AW. 2021. Endogenous and borrowed proteolytic activity in *Borrelia*. Microbiol Mol Biol Rev 85: e00217-20. <https://doi.org/10.1128/MMBR.00217-20>.

Copyright © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Jorge L. Benach, jorge.benach@stonybrook.edu, or A. Wali Karzai, wali.karzai@stonybrook.edu.

Published 12 May 2021

INTRODUCTION

The spirochete genus *Borrelia* (family *Spirochaetaceae*) was named after the French biologist Amedee Borrel (1). It comprises two main groups, the agents of Lyme borreliosis (LB) and relapsing fever (RF), that share common characteristics, of which 21 are members of LB group and 29 are in the RF group (2). The species constituting the LB group are transmitted by the hard ticks of the family Ixodidae (3), whereas the species in the RF group are transmitted by the soft ticks of the family Argasidae. There are two exceptions, *Borrelia recurrentis*, which is transmitted by the body louse, and *Borrelia miyamotoi*, an RF agent transmitted by the hard tick *Ixodes scapularis* (4). LB and RF *Borrelia* spp. share a set of genetic and biological characteristics that unify these organisms as a group. Namely, all LB and RF *Borrelia* spp. have a similar morphology and an obligate parasitic life cycle and are transmitted between vertebrate hosts by arthropod vectors.

The *Borrelia* spp. have an outer membrane, an inner membrane, and a layer of peptidoglycan within the periplasm. A unique feature of the spirochetes, not just the *Borrelia*, is the presence of endoflagella (or periplasmic flagella) in the periplasmic space, between the outer membrane and the peptidoglycan. Spirochetes are highly motile (5). The *Borrelia* spp. have small genomes with a single linear chromosome and several linear and circular plasmids, a unique feature among eubacteria (6). Another characteristic of *Borrelia* is the abundance of lipoproteins (some in the outer membrane), glycolipids, and cholesteryl glycolipids arranged in clusters (7). For a thorough compilation of the biology of *Borrelia*, please see the recent monograph (8).

The trafficking of *Borrelia* spp. between their vectors and vertebrate hosts requires significant physiological changes, including remodeling of their proteome and outer membranes (9). This is an important feature for the topic of this review, since much of this essential remodeling requires the action of an array of proteases. The proteolytic arsenal of *Borrelia* is strategically deployed for disposal of proteins no longer required as it moves from one stage to another or as it is transferred from one organism to another. Likewise, the *Borrelia* spp. are systemic organisms that need to break down and move through certain tissues and barriers, and their proteolytic resources facilitate this movement.

This review focuses on the endogenous protease arsenal of *Borrelia*, the unique features of this proteolytic resource, as well as the borrowed proteolytic capacity through the use of mammalian proteases—notably, those of the fibrinolytic system. Although we consider the proteolytic activity of both LB and RF organisms, the literature is significantly more skewed with studies of *Borrelia burgdorferi sensu lato*, the agent of Lyme disease (10, 11). As a guide, we have included below an abridged mechanistic description of the major protease types discussed in this review.

Serine Proteases

Serine proteases are a large and ubiquitous group that use similar mechanisms of action to cleave peptide bonds. In all cases, the side chain hydroxyl of the catalytic serine (Ser) residue is activated to form a nucleophile that attacks the peptide bond. The catalytic serine residue is activated by deprotonation, either in the context of a catalytic triad (Ser, His, and Asp), a mechanism used by the high-temperature requirement proteases (HtrA) and caseinolytic proteases (ClpP), or a catalytic dyad (Ser and Lys), a mechanism used by the Lon proteases, named for the elongated (long) phenotype of cells lacking these key enzymes. Nucleophilic attack by the activated Ser hydroxyl results in cleavage of the peptide bond such that one fragment of the resulting peptide is released whilst the other remains covalently linked to the active site Ser. A water molecule (H₂O) is coordinated and activated to break the covalently linked intermediate, thus releasing the second peptide fragment and regenerating the active site Ser (12–15).

Threonine Proteases

Threonine proteases use the sidechain hydroxyl group of the catalytic threonine (Thr) residue as the nucleophile that attacks the peptide bond of the target polypeptide. However, the mechanism of action of Thr proteases differs from that of the serine proteases in that the Thr hydroxyl group is activated by the α -amino group of the same Thr residue rather than a catalytic dyad or triad. This restriction necessitates that the nucleophilic Thr be located at the N terminus of the enzyme. Nucleophilic attack by the activated Thr hydroxyl results in cleavage of the peptide bond such that one fragment of the resulting peptide is released whilst the other remains covalently attached to active site Thr. A water molecule (H_2O) is brought to the active site and activated to perform the next step of the reaction, thus releasing the covalently linked second fragment and regenerating the active site Thr (13–15).

Cysteine Proteases

Cysteine proteases use the thiol group of the active-site cysteine (Cys) as a nucleophile to catalyze peptide bond cleavage. The Cys residue is typically found in a catalytic triad with histidine and aspartic acid, akin to serine proteases, or a catalytic dyad with a histidine residue. In the latter case, it is proposed that the thiol group of the active-site Cys residue is more acidic than the hydroxyl of serine proteases, which obviates the need for the additional aspartic acid residue. The overall reaction mechanism of cysteine proteases is similar to that of serine proteases in that the active site histidine removes a proton to activate the Cys thiol group which, in turn, acts as the nucleophile that attacks and cleaves the peptide bond (13–15).

Aspartyl Proteases

Aspartyl proteases use two active-site aspartic acid (Asp) residues; one acts as an aspartate (deprotonated version) to coordinate and activate a water molecule as the nucleophile, while the other functions as an aspartic acid (protonated version) to coordinate the carbonyl ($C=O$) group of the bound polypeptide to make it a better electrophile. The activated hydroxyl (OH) attacks the peptide bond and generates two peptide fragments that are released from the protease without the formation of a covalent intermediate (13–15).

Metalloproteases

Metalloproteases coordinate metal ions that facilitate cleavage of peptide bonds. The majority of metalloproteases use zinc, with a few using other metals such as cobalt. Metalloproteases use a triad composed of several different amino acids, such as histidine, aspartate, glutamate, arginine, and lysine, to bind the metal ion which, in turn, coordinates a water molecule as the ultimate nucleophile. One typically conserved motif, His-Glu-X-X-His (HEXXH), forms the active site of some metalloproteases, where the two His residues help coordinate the metal ion, which, together with the Glu, helps position the bound water molecule. The Glu residue activates the bound water, thus generating a nucleophilic hydroxyl (OH) that attacks the peptide bond. Cleavage of the peptide bond yields two fragments, neither of which is covalently linked to the enzyme (13–15).

ENDOGENOUS PROTEASES OF THE *BORRELIA*

Proteolytic Component of *Borrelia*

The list of 26 annotated proteases of *B. burgdorferi* (Table 1) shows that this bacterium is endowed with a diverse proteolytic arsenal. Of note is the presence of signal peptidases I and II (16). Specifically, there are two signal peptidase I proteins and one signal peptidase II (LspA). LspA has a uniquely important role in the physiology of *B. burgdorferi*, as it cleaves the leader peptides of the large number of lipoproteins that are so critical for this organism. The genomes of relapsing fever *Borrelia hermsii*, an American species, and *Borrelia duttonii*, a sub-Saharan African species, also encode the full complements of these annotated proteases (Table 1). With the exception of HtrA

TABLE 1 List of annotated proteases of *Borrelia burgdorferi*^a

Gene	Locus	Protein	Protease family	Molecular function
<i>lepB</i> (1)	BB_0031	LepB signal peptidase I	Serine-type endopeptidase	Cleavage of signal/leader peptides
<i>map</i> 1	BB_0067	MAP 1	Metalloprotease aminopeptidase	Unassigned peptidases (methionyl aminopeptidase I family)
<i>Ap</i> II	BB_0069	Aminopeptidase II	Metalloprotease	Putative peptidases
<i>htrA</i>	BB_0104	HtrA	Serine endopeptidases	Periplasmic serine protease (Deg family)
<i>map</i>	BB_0105	Methionine aminopeptidase	Metalloprotease	Removes the N-terminal methionine from nascent proteins
<i>rseP</i>	BB_0118	RseP	Metalloprotease	Putative sigma E protease
<i>cym1</i>	BB_0228	Cym1	Zinc metalloprotease	Putative presequence peptidase
<i>pepF</i>	BB_0248	PepF	Serine-type peptidase	Oligopeptidase F
<i>lon1</i>	BB_0253	Lon (La) ATP-dependent protease	Serine-type peptidase	ATP-dependent cleavage of peptide bonds with broad specificity; actual substrates in <i>Borrelia</i> are not known
<i>lepB</i> (2)	BB_0263	LepB signal peptidase I	Serine-type endopeptidase	Cleavage of signal/leader peptides
<i>hslU</i>	BB_0295	HslU ATP-dependent chaperone	ClpYQ (HslUV)	ATPase component of the HslUV system; cleavage of peptide bonds with broad specificity
<i>hslV</i>	BB_0296	HslV peptidase	Threonine-type endopeptidase activity	Peptidase component of the HslUV system; cleavage of peptide bonds with broad specificity
<i>cptA</i>	BB_0359	CptA	Serine-type endopeptidase	Putative carboxypeptidase
<i>apeA</i>	BB_0366	ApeA	Putative aspartyl aminopeptidase	Putative aminopeptidase
<i>clpA</i>	BB_0369	ClpA ATP-dependent chaperone		ATP-dependent cleavage of peptide bonds with broad specificity; actual substrates in <i>Borrelia</i> are not known
<i>lspA</i>	BB_0469	LspA signal peptidase II	Aspartic-type endopeptidase	Proteolipoprotein signal peptides, signal peptidase II
<i>pqqL</i>	BB_0536	PqqL	Zinc metalloprotease	Putative insulinase family member
Unassigned	BB_0592		Glutamate-type endopeptidase	Putative prenyl-processing peptidase family I
<i>lon2</i>	BB_0613	Lon 2 (La) ATP-dependent protease	Serine-type endopeptidase	ATP-dependent cleavage of peptide bonds with broad specificity; actual substrates in <i>Borrelia</i> are not known
<i>clpP1</i>	BB_0611	ClpP peptidase	Serine-type endopeptidase	ATPase component of the ClpXP system; cleavage of peptide bonds with broad specificity; Actual substrates in <i>Borrelia</i> are not known
<i>clpX</i>	BB_0612	ClpX ATP-dependent chaperone		Peptidase component of the ClpX/A/C systems; cleavage of peptide bonds with broad specificity
<i>pepX</i>	BB_0628	PepX/ApeB	Aspartyl-aminopeptidase	Aminopeptidase family member
<i>clpP2</i>	BB_0757	ClpP2 peptidase	Serine-type endopeptidase	Peptidase component of the ClpX/A/C systems; cleavage of peptide bonds with broad specificity
<i>prp</i>	BB_0779	Prp	Cysteine-type endopeptidase	Ribosomal-processing cysteine protease
<i>ftsH</i>	BB_0819	FtsH ATP-dependent protease	Zinc metalloprotease	Degrades misassembled membrane proteins and cytoplasmic regulatory proteins, including σ^{32} , LpxC, and λ CII
<i>clpC</i>	BB_0834	ClpC ATP-dependent chaperone		ATP-dependent cleavage of peptide bonds with broad specificity; actual substrates in <i>Borrelia</i> are not known

^aThe genomes of relapsing fever *Borrelia* species *B. hermsii* and *B. duttonii* carry annotated copies of the genes listed in this table.

(see below), the protease component of all three *Borrelia* species examined is similar. Relapsing fever (RF) and Lyme disease (LD) *Borrelia* spp. are closely related spirochetes but produce different clinical outcomes in animals and humans based on their respective abilities to cause a recurrent spirochetemia or an invasion of multiple organs. Based on our examination of the annotated proteases of the three representative species, it is not possible to conclude that their proteolytic components could contribute to these differences.

The first attempts at detecting proteolytic activity in *B. burgdorferi* utilized collagen (II and IV) and gelatin as the substrates. Collagenolytic activity was detected using spirochete lysates (17). Since then, other proteases have been studied in greater detail. Not surprisingly, despite its relatively small genome size, *B. burgdorferi* has dedicated a larger percentage of its genome to proteolysis than bacteria with larger genomes (Table 2). That *B. burgdorferi* has a full complement of the ATP-dependent proteases (Table 2) indicates that energy-driven proteolysis is a very important physiological feature of this dual-life-cycle bacterium.

The energy-dependent AAA+ (ATPases associated with diverse cellular activities) proteases degrade damaged, misfolded, and surplus proteins and remove key regulatory proteins following major life cycle transitions and growth condition changes. The ATP-dependent proteases that drive these processes must be highly specific to avoid untimely degradation of essential proteins. *B. burgdorferi* contains six ATP-dependent proteases: ClpXP, ClpAP, ClpCP, HslUV, FtsH, and Lon (Tables 1 and 2).

Energy-dependent proteases are composed of an AAA+ component and a peptidase component. These two components can be encoded by the same gene, as is the case for Lon (Fig. 1, left) and for FtsH, or by two distinct genes, one encoding the AAA+ chaperone/unfoldase (ClpX, ClpA, and ClpC) and the other encoding the ClpP peptidases (Fig. 1, right). Furthermore, the unfoldases belong to the Hsp100/Clp family of proteins and contain one (in the case of ClpX) or two (in the case of ClpA, ClpB, or ClpC) ATPase domains. All of these ATP-fueled nanomachines can unfold and degrade folded proteins. The active AAA+ protease form ring-shaped oligomeric assemblies (Fig. 1) with a central pore leading to the sequestered peptidase chamber (18). While most Gram-negative bacteria contain ClpXP and ClpAP, Gram-positive bacteria typically contain some combination of ClpXP, ClpCP, and ClpEP. For instance, *Staphylococcus aureus* contains ClpXP and ClpCP, where ClpCP is more effective at degrading damaged proteins than ClpXP (19). Curiously, *B. burgdorferi* has all three Clp proteases (ClpAP, ClpXP, and ClpCP) and carries two ClpP peptidases, ClpP1 and ClpP2 (Table 2).

HslUV, also known as ClpYQ, is formed by a union of the dodecameric peptidase HslV and its hexameric, ATP-dependent partner, HslU. HslUV is an AAA+ protease found in certain bacteria, including *B. burgdorferi* (Table 2), and in the mitochondria of some lower eukaryotes, such as trypanosomatids, and shares sequence, structural, and mechanistic similarities with the more sophisticated 20S proteasome. The HslU unfoldase regulates the proteolytic activity of the HslV peptidase and is responsible for the substrate specificity of the protease (20, 21). Moreover, HslUV and ClpXP have been implicated in the regulation of the SOS response, polysaccharide synthesis, RNA metabolism, and symbiotic nitrogen fixation (22). The FtsH and Lon proteases of *B. burgdorferi* are considered below.

Amino and Carboxypeptidases of *B. burgdorferi*

An aminopeptidase from *B. burgdorferi* was shown to have aminopeptidolytic activity in spirochete lysates. The purified protein is a 300-kDa hexamer formed by 50-kDa monomers. The enzyme was identified as zinc-dependent aminopeptidase II (BB0069) with significant identity to the M29/T family of metallopeptidases (23).

CtpA, a carboxyl-terminal protease in *B. burgdorferi* is involved in the processing of P13 and BB0323, and *ctpA* inactivation has a pleiotropic effect on *Borrelia* protein synthesis (24). Site-directed mutagenesis to alter CtpA cleavage sites of the outer membrane (OM) porin P13 resulted in ectopic expression of a C-terminally truncated P13.

TABLE 2 ATP-dependent proteases of selected bacterial and other species

Species	No. of encoded proteins	Lon	ClpX	ClpP	ClpA	ClpC	FtsH	HslUV
<i>E. coli</i>	4289	✓	✓	✓	✓	✓	✓	✓
<i>B. subtilis</i>	4100	✓	✓	✓	✓	✓	✓	✓
<i>Mycobacterium tuberculosis</i>	3918	✓	✓	✓ (ClpP1 and ClpP2)	✓	✓	✓	✓
<i>Synechocystis</i>	3169	✓	✓	✓	✓	✓	✓	×
<i>Deinococcus radiodurans</i>	2580	✓	✓	✓	✓	✓	✓	×
<i>Thermotoga maritima</i>	1846	✓	✓	✓	✓	✓	✓	✓
<i>H. influenzae</i>	1709	✓	✓	✓	×	×	✓	✓
<i>Aquifex aeolicus</i>	1522	✓	✓	✓	✓	✓	✓	✓
<i>H. pylori</i> 26695	1566	✓	✓	✓	✓	×	✓	✓
<i>Treponema pallidum</i>	1031	✓	✓	✓	✓	✓	✓	?
<i>Chlamydia pneumoniae</i>	1052	✓	✓	✓ (ClpP1 and ClpP2)	✓	✓	✓	×
<i>Chlamydia trachomatis</i>	894	✓	✓	✓ (ClpP1 and ClpP2)	✓	✓	✓	×
<i>B. burgdorferi</i>	850	✓ (Lon1 and Lon2)	✓	✓ (ClpP1 and ClpP2)	✓	✓	✓	✓
<i>B. hermsii</i>	827	✓ (Lon1 and Lon2)	✓	✓ (ClpP1 and ClpP2)	✓	✓	✓	✓
<i>B. duttonii</i>	850	✓ (Lon1 and Lon2)	✓	✓ (ClpP1 and ClpP2)	✓	✓	✓	✓
<i>Rickettsia prowazekii</i>	834	✓	✓	✓	×	×	✓	✓
<i>Mycoplasma pneumoniae</i>	677	✓	×	×	×	×	✓	×
<i>Mycoplasma genitalium</i>	467	✓	×	×	×	×	✓	×
<i>N. gonorrhoeae</i>	2030	✓	✓	✓	✓	×	✓	?
<i>Caulobacter crescentus</i>	3884	✓	✓	✓	✓	×	✓	✓
<i>Y. pestis</i>	3873	✓	✓	✓	✓	×	✓	✓
<i>Streptococcus pyogenes</i>	1680	✓	✓	✓	×	✓	✓	?
<i>Staphylococcus epidermidis</i>	2326	✓	✓	✓	×	✓	✓	✓
<i>Listeria monocytogenes</i>	2930	✓	✓	✓	✓	✓	✓	✓
<i>Agrobacterium tumefaciens</i>	5163	✓	✓	✓	✓	×	✓	✓
<i>Clostridium tetani</i>	2692	✓	✓	✓	✓	✓	✓	?
<i>Arabidopsis thaliana</i>	29085	✓	✓	✓	✓ limited homology	✓	✓	✓
<i>Homo sapiens</i>	~21000	✓	✓	✓	✓ limited homology	×	✓	×

Additionally, a lower-molecular-weight variant of surface lipoprotein OspC was mislocalized to the periplasm. Further investigation revealed that the OspC variant resulted from C-terminal proteolysis by CtpA. These findings emphasized that the CtpA of *B. burgdorferi* is not specific for targeting proteins that lack structurally constrained C termini, as proteolysis appears to occur independently of a specific peptide recognition sequence (25).

FtsH of *B. burgdorferi*

Molecular chaperones ensure correct protein function by supervising protein folding and shuttling proteins to their final destinations. Misfolded, aberrant, and unwanted proteins, on the other hand, are targeted for proteolytic degradation. The membrane-anchored, 71-kDa ATP-dependent zinc metalloprotease FtsH fulfills these functions in bacteria, mitochondria, and chloroplasts, where it degrades cytoplasmic and membrane proteins. While FtsH is essential for the viability of *Escherichia coli*, *Bradyrhizobium japonicum*, and *Helicobacter pylori*, in other bacteria, it is important for cell viability only during stress conditions and stationary-phase growth. Known FtsH substrates include the heat shock response sigma factor σ^{32} and the lipid A biosynthesis enzyme LpxC. In fact, absence of FtsH perturbs the balance of lipopolysaccharide and phospholipid synthesis, resulting in cell death. The SecYEG translocon, which facilitates protein transport across the inner membrane, is also subject to proteolytic regulation by FtsH. When not in complex with SecE and SecG, SecY is recognized and degraded by FtsH. Additionally, FtsH degrades KdtA, a transferase required for the biosynthesis of oligosaccharides (26, 27).

The FtsH of *B. burgdorferi* (FtsHBb) has the ATP-binding consensus sequences and the conserved proteolytic HEAGH (HEXXH) motif, suggesting it functions like the canonical FtsH from *E. coli*. FtsH degrades misfolded cytosolic and membrane proteins and possess the capacity to function as a chaperone (28). The canonical *E. coli* FtsH




FIG 1 Architectural arrangements of AAA+ proteases. (Left) The structural model of Lon protease hexamer in the active substrate-engaged state is shown (PDB 6ON2). Each Lon monomer contains both the ATPase and the peptidase domains. In this depiction, one of the six Lon protomers is omitted to highlight the bound substrate (yellow), staircase arrangement of the active substrate engaged enzyme, and the peptidase active-site Ser-Lys dyad (purple). (Right) The structural model of the ClpAP protease in the active substrate-engaged state is shown (PDB 6W1Z). The ClpA ATPase forms a homohexamer that sets on top of the assembled ClpP peptidase, which is composed of two stacked heptamers (dark red and red, respectively). In this depiction, one of the six ClpA protomers is omitted to highlight the two AAA+ domains and the bound substrate (yellow).

associates with modulators HflC/HflK for oligomerization (28–31). The presence of the three proteins (HflC, HflK, and FtsH) in lipid rafts of *B. burgdorferi* strongly indicates that the complex carries out its activity in these microdomains (32, 33). The genes for the FtsH protease and its modulators, HflK and HflC, were deleted, with interesting and unexpected results. FtsH depletion (but not overproduction) in *B. burgdorferi* resulted in membrane deformation and cell death. In contrast, deletion of the FtsH modulators HflK and HflC (Δ HflK/C) did not alter morphology, growth rate, growth under stress conditions, or infectivity (26).

Lon Proteases of *B. burgdorferi* Are Unusual in the Eubacteria

As a member of the AAA+ protease family, Lon can be found in bacteria, archaea, and eukaryotes. Extensively studied in the context of bacterial proteolysis, the ATP-dependent Lon protease regulates a variety of cellular processes, such as capsule synthesis, genetic competence, cell motility, cell division, DNA replication, heat shock response, and pathogenesis. Lon is also thought to be responsible for degrading approximately 50% of all misfolded proteins in *E. coli*, highlighting its integral role in preserving cell viability. As a proteolytic regulator, Lon degrades various native regulatory proteins that modulate major cellular processes. These include the following: HU β , the DNA-binding histone-like protein that plays a role in DNA replication and gene regulation; Sula, a critical cell division protein; RcsA, the capsule synthesis regulator; CcdA, an antitoxin that is part of the F plasmid-based killing system; IbpA, a heat shock protein; SoxS, a transcription activator that plays an important role in removing reactive oxygen species; SwrA, the master regulator of flagellar biosynthesis and cell motility; and transfer-messenger RNA (tmRNA)-tagged proteins (34–40). A common feature among proteases, Lon is assisted by adaptor proteins (41) that enhance its activity and fine tune its substrate specificity. Two such Lon-specific adaptors have been identified: SmiA, which mediates the degradation of SwrA in *Bacillus subtilis*, and HspQ (39, 42). As one of the most intriguing adaptor molecules, HspQ is itself a Lon substrate while also acting as an allosteric activator of the protease. Lon-mediated engagement and degradation of HspQ enhances the proteolysis of several native regulatory substrates, such as YmoA (*Yersinia* modulating protein A), Y0390, Fur (ferric uptake regulator), and RsuA (ribosomal small subunit pseudouridine synthase A) (42, 43). Lon protease of *Salmonella* degrades the heat-stable nucleoid structuring protein (H-NS),




FIG 2 Architectural arrangements of HtrA and DegP proteases. (Left) The structural model of HtrA protease trimer in the active peptide-bound state is shown (PDB 2Z9I). Each HtrA monomer is shown in a different color, with two monomers containing a bound autoproteolytic peptide (yellow) to the active site. The third monomer is shown without the bound peptide to highlight the active-site Ser-His-Glu catalytic triad (red). (Right) A structural model of the DegP protease dodecamer in the substrate-bound state (PDB 4A8D) is shown. The DegP chaperone forms a cage-like structure composed of 4 trimers, each depicted in a different color, with the bound substrate (outer membrane protein [OMP]; yellow) captured inside the dodecameric cage.

unleashing a complex chain of events that enables the pathogen to express foreign virulence genes during infection (43).

B. burgdorferi harbors two Lon proteases, Lon-1 and Lon-2 (44), which is a rare but not unique occurrence. The first evidence for the presence of the *lon* gene of *B. burgdorferi* yielded a protein of 806 amino acids with substantial sequence identity to other bacterial Lon proteases. Transcriptional upregulation of *lon-1* follows exposure to blood *in vitro* (45). The *lon-2* gene is immediately downstream of ATP-dependent proteases *clpP* and *clpX*, as is in *E. coli lon*. Lon-1 and Lon-2 of *B. burgdorferi* cluster separately in the N-terminal substrate-binding domains, and this could reflect differences in substrate specificity. Recombinant Lon-1 had the properties of an ATP-dependent chaperone protease, with a catalytically active serine-lysine dyad, that has caseinolytic activity *in vitro*. However, the recombinant Lon-1 could not degrade an ssrA-tagged substrate (41, 46). Moreover, *lon-1* did not complement an *E. coli lon* mutant, while Lon-2 corrected two characteristic phenotypes of the *lon*-mutant. Lon-2 functions in a manner consistent with canonical Lon, engaged in cellular homeostasis. Thus, Lon-1, as a result of its blood induction and as a unique feature of *Borrelia*, is important during the transition from the arthropod vector to the mammalian host.

The role of the two *Borrelia* Lon proteases in murine infection indicate that Lon-1 plays a critical role for the infection of *B. burgdorferi* (47). A *lon-1* deletion mutant was attenuated in mice and displayed other characteristics such as growth defects in BSK-II medium and resistance to osmotic stress. Production of BosR, RpoS, and OspC increased in the mutant, suggesting that one or all of these regulatory proteins may be Lon-1 substrates. A catalytic-site Ser-to-Ala mutant did not infect mice, suggesting that the proteolytic activity of Lon-1 is essential for infection (47, 48). A *lon-2* deletion mutant was attenuated in mice but did not have a growth defect in culture (48). This mutant showed resistance to osmotic stress, and protein levels of RpoS and OspC were decreased. The finding that neither *lon1* nor *lon2* is essential may indicate that their activities are synergistic and that, despite their biochemical differences, one can compensate, at least partially, in the absence of the other.

HtrA of *Borrelia*, a New Strategy for Pathogenesis

The HtrA family of serine proteases can be found in all cells, from prokaryotes to primates. Unifying features of this family are the trimeric structure and the proteolytic domain catalytic triad composed of Ser-His-Asp (Fig. 2). These proteases also have one or two C-terminal PDZ domains that mediate protein-protein interactions (49).

E. coli DegP was the first periplasmic HtrA protease characterized (50, 51). It functions as a protease to degrade misfolded proteins and as a chaperone during the protein-folding stress response (51). As a chaperone, DegP protects proteins from

degradation in the periplasm, and as a protease, it degrades selected substrates (52). The DegP trimer can auto-oligomerize after binding a misfolded or partially folded substrate, forming a macromolecular cage-like structure (Fig. 2B) that functions as a chaperone to protect the trafficking of outer membrane proteins through the periplasm (53, 54).

The first identification of the HtrA homolog (BhpA) in the genus *Borrelia* was made in two species of relapsing fever spirochetes, *B. hermsii* and *B. turicatae*. Although most relapsing fever organisms have the *bhpA* gene, it is not shared with *B. burgdorferi* (55). Recombinant BhpA protein degrades β -casein and is transcribed at all growth temperatures *in vitro* but at much higher levels during *B. hermsii* infection in mice. A gene encoding BtpA, an HtrA family protease, is present in *B. turicatae* (55). Transcriptional analyses in *B. turicatae* revealed that *btpA* was expressed as part of an operon. Indeed, it was demonstrated that BtpA and proteins encoded by two adjacent genes were produced in response to culture at mammalian body temperature. Mutants of *btpA* exhibit no growth defects in response to heat shock but are more sensitive to oxidative stress. The fact that the *btpA* mutant is infectious in the murine model indicates that its function is not required for pathogenesis (56). A phylogenetic analysis showed that *B. hermsii* BhpA is the homolog of BhtA of *B. turicatae* but not of *B. burgdorferi* HtrA (48). Our own phylogenetic analysis agreed with these results (50).

B. burgdorferi harbors a gene for a single HtrA protease (HtrABb) on its chromosome that shares ~40% amino acid homology with DegP from *E. coli*. This protease has been the subject of several studies that disclosed some unique features as well as a possible role in pathogenesis. HtrABb processes a conserved protein substrate, BB0323, into N- and C-terminal fragments that are biologically active and supports spirochete growth (57, 58). The cleavage of BB0323 has critical functional implications in the spirochetal life cycle, as the N-terminal fragment is important for cell fission and the C-terminal LysM domain fragment is essential for mammalian infection. Biochemical studies showed that HtrABb, like DegP, has the trimer as its fundamental unit (Fig. 2A). Recombinant HtrABb degrades casein, while its catalytic serine mutant (HtrABbS198A) does not (59). Despite its homology to DegP, HtrABb does not complement an *E. coli* *degP* deletion mutant. Two additional HtrABb substrates, basic membrane protein D (BmpD/BB0385) and chemotaxis signal transduction phosphatase CheX (BB0671), have been identified. BmpD is an important immunogenic adhesin expressed in infected patients, and CheX is a crucial protein for motility (59). HtrABb may function in regulating outer membrane lipoproteins and in modulating the chemotactic response of *B. burgdorferi*. Proteolytic activity of HtrABb is inhibited by micromolar concentrations of zinc preferentially over copper and manganese (60). FlhD, a flagellar cap protein, promotes flagellin (FlaB) polymerization and filament growth within the periplasm. Deletion of *flhD* leads to an accumulation of unpolymerized FlaB, which is degraded by HtrA in the periplasm (61). HtrABb-dependent proteolysis of FlaB and CheX, as well as the defect on swarm assays, suggests that HtrABb plays an important role in regulating *B. burgdorferi* motility (5, 61).

An HtrABb overexpression approach identified outer membrane protein P66 as a substrate. Both P66 and HtrA partitioned into detergent-resistant membranes, which contain cholesterol-glycolipid-rich membrane regions known as lipid rafts. This agreed with previous work which showed that HtrABb and p66 are constituents of *B. burgdorferi* outer membrane vesicles (62). Their colocalization establishes that they could interact efficiently, and their protease/substrate relationship provides functional relevance to this interaction. The overexpresser strain showed reduced levels of *p66* transcript, indicating that HtrABb-mediated regulation of *p66* may occur at multiple levels. There is increasing evidence that HtrABb could be located in a functional form outside the periplasm (63). An HtrABb-overexpressing strain of *B. burgdorferi* (A3HtrAOE) revealed that protein levels of P66 were reduced compared to that in wild-type *B. burgdorferi*, consistent with its being a substrate of this protease. In addition, Hbb, a *p66*-DNA-binding transcription factor (64), was specifically degraded by HtrABb, providing further

evidence for a role for both in the regulation of P66. P66 is a very important outer surface protein for *B. burgdorferi*, as it functions as both an adhesin and a porin (65–67).

The A3HtrAOE strain displayed reduced motility in swarm assays, indicating a link between overexpression of HtrABb and its enzymatic specificity for P66. However, the $\Delta p66$ strain did not have reduced motility in the swarm assays, negating a role for this protein. The proteomics analyses also showed that overexpression of HtrABb had an impact in the production of the cellular levels of three glycolytic enzymes, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), glycerol-3-phosphate dehydrogenase (GPDH), and glycerol kinase (GK), and an enzyme involved in carbohydrate metabolism, diphosphate-fructose-6-phosphate 1-phosphotransferase. Consistent with the reduced protein levels of these glycolytic enzymes, A3HtrAOE was also deficient in production of pyruvate. These experiments suggested a role for HtrABb in regulating the metabolic activity of *B. burgdorferi* (68).

Lmp1, a surface-exposed protein involved in various functions in spirochete infectivity, is another HtrABb substrate (69). HtrABb degrades Lmp1 into polypeptide fragments that are essential for transmission of *B. burgdorferi* from ticks to mammals (69). That Lyme disease patients, as well as mice experimentally infected with *B. burgdorferi*, produce antibodies to HtrABb (59, 70, 71) prompted an attempt to determine whether recombinant HtrABb could be used as a vaccine, but it failed to protect the mice after a challenge with the organisms (72).

Recent studies succeeded in creating a targeted deletion of HtrABb, suggesting a nonessential role for this protease in microbial viability. However, the mutant displayed growth, morphological, and structural defects during cultivation at 37°C, confirming a high-temperature requirement for protease activation and function. Moreover, HtrABb-deficient spirochetes were unable to establish infection in mice. Earlier studies confirmed HtrA-mediated proteolytic processing of BB0323 (discussed above), the lack of which likely contributed to the inability of the mutant strain to survive in a mammalian host (73).

Canonical HtrA is a periplasmic protease, but there is enough evidence to support proteolytic activity in other bacterial cell compartments as well as in outer membranes. The location of HtrABb outside the canonical periplasm was established through proteinase K studies that demonstrated its surface exposure. Using sera from patients with early and late Lyme disease, this study also established that HtrABb is immunogenic, thus confirming its expression during the disease process. Additional HtrABb extracellular substrates have been identified in the tissues preferred by *B. burgdorferi*. Aggrecan, a cartilage-specific proteoglycan, is an HtrABb substrate (71) as are proteoglycans of the extracellular matrix (ECM) and fibronectin. Fragments derived from the degradation of fibronectin amplify the inflammatory process, thus contributing to the pathogenesis of Lyme disease. Exposure of chondrocytes to recombinant HtrABb leads to the production of proinflammatory cytokines and chemokines that are well-known participants in the Lyme arthritis milieu. Therefore, HtrABb is thought to actively participate in dissemination and in tissue damage due to inflammation triggered by Lyme disease (70).

We found that HtrABb was one of a group of proteins detected by mass spectrometry in isolated *B. burgdorferi* vesicles (32, 33). HtrABb exists in both membrane-bound and soluble forms and is detectable in conditioned medium (59, 70). Indeed, exo-periplasmic location and activity for HtrA have been well established in other pathogens such as *Campylobacter jejuni* (74) and for extracellular secretion and degradation of E-cadherin by *Helicobacter pylori* (75, 76). The same is true for secreted HtrA proteases from other Gram-negative pathogens in cleaving junctional proteins (75, 76). Extracellular proteolysis by secreted HtrA is thought to augment the invasiveness of pathogenic bacteria by degrading host ECM components and cellular junctional molecules. This topic is considered in a review where the proteolytic activities of secreted HtrA from both Gram-positive and Gram-negative pathogens promote invasiveness and cell damage and thus constitutes a novel mechanism for pathogenesis (77). It is




FIG 3 Mammalian plasminogen activation system indicating all of its components that interact with *Borrelia* species.

proposed that HtrA export is a newly acquired strategy used by an increasing number of bacterial pathogens.

BORRELIA AND BORROWED PROTEOLYSIS

This section considers the vast literature on the utilization of host proteases by *Borrelia*. Research into borrowed proteolysis has yielded valuable insights into spirochetal biology.

Plasminogen Activation System

The plasminogen activation system (PAS) is a highly controlled system for dissolving fibrin clots (thrombi) that has been implicated in a diverse set of normal and pathological processes. Its central effector element, plasminogen (PLG), is a zymogen initially expressed in humans as an 810-amino-acid polypeptide that is subsequently modified to a 791-amino-acid mature protein (78). PLG circulates in the blood and tissue fluids at a concentration of approximately $2\ \mu\text{M}$ (79) and is activated enzymatically to form active plasmin through cleavage of its Arg⁵⁶¹-Val⁵⁶² bond. Plasmin has a broad trypsin-like range of specificity, enabling it to degrade a wide variety of substrates in addition to its canonical role in thrombolysis (Fig. 3).

This powerful source of proteolysis, due to its high concentration in circulation, is tightly controlled by a set of two PLG activators (PAs), originally identified in urine and tissue extracts and that have been classified as belonging to distinct classes. Urokinase-type plasminogen activator (uPA) is a 411-amino-acid zymogen (pro-uPA) that is activated by plasmin cleavage at Lys¹⁵⁸-Ile¹⁵⁹ to generate uPA, a two-chain enzyme connected by a disulfide bond. Tissue-type plasminogen activator (tPA) is a proteolytically active, 527-amino-acid single-chain molecule that is also processed by plasmin into a two-chain form, which is substantially more active than the single-chain form. uPA and tPA are alike in that they are activated by plasmin but have mainly divergent roles, with uPA being linked to inflammation, wound healing, tissue remodeling, metastasis, and invasion (80–82).

Unchecked plasmin elicits a cascade of proteolytic events that can lead to degradation of ECM, diseases of the vasculature, and cancer metastasis (83). To prevent uncontrolled plasmin activity and pathological tissue damage, a system of protease inhibitors, acting either at the plasmin level or at the PA level, is constitutively expressed. At the plasmin level, alpha2-antiplasmin (α 2-AP), a 70-kDa serpin, inhibits plasmin very rapidly, forming a 1:1 stoichiometric complex. At the PA level, serpin plasminogen activator inhibitor (PAI-1) and, to a lesser extent, PAI-2, PAI-3, and protease nexin inhibit uPA and thus plasmin, by binding to the uPA active site (84, 85).

Urokinase plasminogen activator receptor (uPAR; CD87), a glycosylphosphatidylinositol (GPI)-anchored 55- to 60-kDa protein (86–89) first identified in monocytes (90, 91), binds to uPA, pro-uPA, and the ECM glycoprotein vitronectin (92). Plasmin bound to uPAR is protected from inactivation by plasminogen activator inhibitors (PAI) and α 2-AP (unbound plasmin is efficiently inactivated by PAIs). In addition, plasmin bound to fibrin is several orders of magnitude more resistant to inactivation by α 2-AP (78). In both normal and pathological settings, cell-associated plasmin acquisition is believed to be a product of cell-associated uPA activation of PLG. A keynote occurrence in this process is the interaction between uPA and its high-affinity receptor, uPAR (86, 87, 89). The focus of uPA and, subsequently, plasmin generation on the cellular leading edge is thought to be the means by which inflammatory and cancer cells (uPAR is overexpressed in metastatic cancers of various types) degrade ECM and migrate from one anatomical site to another. This supports its role as a central player in mediating ECM degradation during inflammation and cancer progression (93).

Plasmin(ogen) Binding and Invasion by Bacteria

In addition to innate genome-encoded proteases, there is an expanding collection of bacteria (both Gram positive and Gram negative) that acquire proteolytic activity by borrowing host PAS components. This occurs through binding of five PLG kringle structures to exposed lysine residues on the bacterial surface (94, 95). Once bound, the PLG can be activated to plasmin, either by native PAs or by further interaction with the host PAS through binding of uPA or, less commonly, tPA. The binding of plasmin to lysine residues on the bacterial surface is inhibited by competition with the lysine analogs ϵ -aminocaproic acid (EACA) and tranexamic acid, confirming its lysine specificity (96). At the plasmin level, lysine binding is inhibited by the presence of the physiological inhibitor α 2-AP; however, bound plasmin is protected from deactivation by α 2-AP (79, 97–99). Supplementary plasmin activity inhibition can also be achieved by nonspecific plasma serpin α 2-macroglobulin (100). Interestingly, the iron-binding milk lactoferrin blocks PLG conversion to plasmin by binding to PLG (101).

PLG-binding bacteria can be divided into two basic categories: those that express endogenous PAs and those that do not. Among those that produce their own PAs are *Streptococcus pyogenes*, *Staphylococcus aureus*, and *Yersinia pestis* (streptokinase, staphylokinase, and plasminogen activator Pla, respectively) (102–107). Streptokinase and staphylokinase are not proteases but form a 1:1 complex with PLG and plasmin, exposing the active site of plasmin and causing conformational changes leading to increased specificity and efficiency for both (41, 108, 109). In contrast to streptokinase and staphylokinase, Pla is a protease and functions in a manner similar to that of uPA and tPA by activating PLG-to-plasmin conversion through proteolytic cleavage of the Arg⁵⁶⁰-Val⁵⁶¹ peptide bond (110). The bacteria that borrow PAs from the host in order to activate bound PLG include *Haemophilus influenzae*, *Moraxella catarrhalis*, *Proteus mirabilis*, *Pseudomonas aeruginosa* (111, 112), *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Helicobacter pylori* (113, 114), *Salmonella enterica* serovar Typhimurium (110), *Escherichia coli* (115, 116), and *Borrelia* spirochetes, including *B. burgdorferi sensu stricto* (3, 117), *B. garinii*, *B. afzelii* (*sensu lato*), *B. recurrentis* (118), *B. hermsii* (119), and other *Borrelia* spp. (120).

Plasmin(ogen) Binding by *Borrelia*

The expanding literature on *Borrelia* and the PAS justifies its treatment as a subject on its own. For that reason, the focus of this review section will be on the interaction with the host PAS by *B. burgdorferi* and related *Borrelia* species only; for reviews of other bacteria and PLG, see reference 110. Figure 3 shows the high level of utilization of the PAS by *Borrelia*. It can be appreciated that all the components of the PAS can interact with these organisms. The affinity of *Borrelia* for PLG is firmly grounded on the unique aspects of its biology. The genome of *B. burgdorferi* is A+T rich (72%) (6), as are the genomes of two RF species, *B. parkeri* (121) and *B. hermsii* (122). The most frequently used codon is AAA, which, along with AAU, codes for lysine, the amino acid

associated with binding to PLG. Not surprisingly, the lysine content of *B. burgdorferi* is 10.2% (6). The ubiquity of lysine residues in *Borrelia* is not only the major factor influencing binding of these organisms to PLG but is also contributory to binding to negatively charged molecules such as proteoglycans and others. Although outside this review, the genomes of *Staphylococcus aureus* and species of *Streptococcus*, also notable for PLG binding, are similarly A+T rich.

B. burgdorferi, *B. garinii*, *B. afzelii*, *B. spielmanii*, and *B. miyamotoi*, the major agents of Lyme disease and miyamotoi disease in North America, Europe (117, 123–126), and Asia (4, 127–130), are transmitted by hard ticks of the genus *Ixodes*. These pathogens spread hematogenously from the initial site of tick attachment to colonize a variety of tissues and organs, causing joint swelling, complications of the heart and nervous systems, and acrodermatitis chronica atrophicans, a late skin manifestation (131–133). The penetration of tissue barriers (capillary endothelium, ECM, and basement membranes) by the spirochete is a requirement to establish systemic Lyme disease infection and necessitates the proteolytic breakdown of these aforementioned ECM components (134, 135).

Although *Borrelia* spp. have been shown to adhere to a large diversity of mammalian and tick cells, mostly under laboratory conditions (136), the preferred site, as observed *in vivo*, is in the ECM and connective tissue (137–143).

In addition to *B. burgdorferi* (*sensu stricto*), the binding of PLG to spirochetal outer membrane has been documented to occur in a variety of other *Borrelia* species, including agents of Lyme disease in Europe, *B. garinii* and *B. afzelii* (*sensu lato*), and in North America/Asia (144, 145). Relapsing fever *Borrelia* species *B. hermsii* (119, 146), *B. coriaceae*, *B. parkeri*, *B. turicatae* (144), and *B. crocidurae* (147) also have been shown to bind plasmin(ogen). In addition to that of PLG, binding of uPA to the outer membrane has also been documented in *B. burgdorferi* as well as in other *Borrelia* species. The binding was diffuse in species other than *B. burgdorferi* but was localized to the spirochete terminus. In mice, uPA-coated *B. burgdorferi* exhibited increased infectivity (144, 148). The PAS also has a role in heart and brain invasion by relapsing fever *Borrelia*, resulting in organ injury (143, 147). RF spirochetes utilize the PAS for organ invasion, as has been shown in *in vivo* studies.

Substrates Degraded by *Borrelia* with Bound Plasmin

Degradation of chromogenic substrate S-2251 has been utilized to identify plasmin-bound proteolytic activity in bacterial species, including *Borrelia* (142, 149, 150). The plasmin in turn can degrade glycoprotein ECM components such as fibronectin (149), laminin, and vitronectin but not collagen (148, 151). *B. burgdorferi* complexed with plasmin was shown to degrade an *in vitro* ECM model (Table 3) (142). Spirochetes with bound PLG or uPA alone as well as spirochetes with bound plasmin in the presence of serine protease inhibitor aprotinin did not show any significant ECM proteolysis (151, 152). Such a process occurring *in vivo* would represent a powerful mechanism for the spirochete to disseminate from the bloodstream to colonize distant tissues/organ sites in the host.

Borrelia and Plasminogen Receptors

The localization of plasmin(ogen) to the *B. burgdorferi* outer membrane has led to investigations to identify receptor proteins accessible for PLG binding. Among the many reported *B. burgdorferi* PLG-binding proteins, several have been characterized (Table 3). Outer membrane lipoprotein OspA was implicated early on as a spirochetal receptor for PLG (142, 149, 152). It is now accepted that OspA, under the control of alternative sigma factor RpoS (153), is downregulated soon after *B. burgdorferi* is transmitted to the mammalian host (154), throwing into question its biological relevance for plasmin-mediated pathogenesis in mammals. OspA may still play an important role in PLG-mediated spirochete dissemination in the tick (155). On the other hand, outer membrane lipoprotein OspC, a 21-kDa lipoprotein critical for infection in mice (156), has been proposed as a biologically relevant PLG receptor (157, 158). In strains where it is highly expressed, OspC binds PLG, and the two molecules colocalize in the outer membrane, which facilitates its

TABLE 3 *Borrelia* ligands for plasminogen

<i>Borrelia</i> PLG ligand	Species	Disease	PLG ligand (kDa)	Reference(s)
OspA ^a	<i>B. burgdorferi</i>	LD ^b	30	142, 149
OspC	<i>B. burgdorferi</i>	LD	21	158
Enolase	<i>B. burgdorferi</i>	LD	47	170, 171, 172
BpcA	<i>B. parkeri</i> , <i>B. turicatae</i>	RF ^c	17	164
ErpP ^d (CRASP ^e -3)	<i>B. burgdorferi</i>	LD	19	159
ErpC (CRASP-4)	<i>B. burgdorferi</i>	LD	60	159
ErpA (CRASP-5)	<i>B. burgdorferi</i>	LD	18	159
CRASP-1 (CspA)	<i>B. burgdorferi</i> , <i>B. afzelii</i> , <i>B. spielmanii</i>	LD	28	120, 161, 192
BhCRASP1/FH	<i>B. hermsii</i>	RF	21.5	163
BBA70	<i>B. burgdorferi</i>	LD	~30	150
CbiA	<i>B. miyamotoi</i>	MD ^f	21	145
70-kDa BPBP	<i>B. burgdorferi</i>	LD	70	152, 175
HcpA	<i>B. recurrentis</i>	RF		118
FhbA ^g	<i>B. hermsii</i>	RF	20.5	119

^aOspA, outer surface protein A.^bLD, Lyme disease.^cRF, relapsing fever.^dErp, OspF/E-related protein.^eCRASP, complement regulator-acquiring surface protein.^fMD, miyamotoi disease.^gFhbA, factor H binding protein A.

conversion to proteolytically active plasmin with the addition of exogenous uPA. This colocalization is not observed in an OspC knockout (158). The lack of significant PLG binding in an OspC-deficient *B. burgdorferi* strain is seemingly in contrast with the literature, since there has been evidence given for alternative PLG-binding sites in several studies. For example, outer membrane ErpP proteins (OspE/F-related), also known as complement regulator-acquiring surface protein (CRASP)-3, CRASP-4 (ErpC), and CRASP-5 (ErpA), have also been reported to bind PLG where it is converted to plasmin, possibly increasing the spirochetes' ability to disseminate through proteolysis of the host ECM (159). Members of the CRASP family also bind complement regulator factor H, FHL-1 (factor H-like protein), and complement factor H-related protein (160, 161), which serve to ward off infection, as well as laminin (159, 162). In another study, CRASP-1 bound PLG and cleaved chromogenic substrate S-2251 (161).

Increasing evidence shows that the affinity for host PLG and components of the complement system is also a unifying characteristic among various *Borrelia* species other than *B. burgdorferi sensu stricto*. *B. afzelii* and *B. spielmanii* (126) as well as *B. burgdorferi* CspA (CRASP-1) bound factor H and PLG, concurrently and once the PLG was activated by uPA, it was able to partially degrade central complement protein C3b (163). Additionally, relapsing fever spirochete *B. hermsii* expresses a novel form of CRASP-1 (designated BhCRASP-1), bound to complement regulators as well as PLG (163). Surface lipoprotein BpcA from *B. parkeri* bound factor H and factor H-related protein 1 as well as PLG, degraded S-2251 and fibrinogen, and conferred resistance to complement-mediated killing. The BpcA homologue from *B. turicatae* failed to bind complement system regulators, while its ability to bind PLG was preserved (164). Surface protein FhbA, a 20.5-kDa lipoprotein from *B. hermsii*, bound factor H and PLG, identifying it as an important molecule in RF pathogenesis (119, 165). Finally, *B. miyamotoi*, the RF agent of miyamotoi disease, a Lyme disease-like syndrome, expresses a surface-exposed protein, CbiA, that exhibits PLG-binding properties, and the bound PLG can be converted to active plasmin capable of cleaving fibrinogen (145).

The enzyme enolase is a well-known component of the glycolytic pathway which catalyzes the conversion of 2-phosphoglycerate to phosphoenolpyruvate. In some bacteria, enolase is also a surface-exposed protein that functions as a PLG receptor (166–169). Accumulating evidence points to a similar role in *B. burgdorferi* (170–172). *B. burgdorferi* enolase, in addition to its cytoplasmic function in glycolysis, would also

have to be exported to the spirochetal outer membrane for it to bind plasmin(ogen) and facilitate extracellular proteolysis needed for spirochetal spread across tissue barriers. Experimental evidence lends credence to this idea. *B. burgdorferi* enolase has been shown, through immunoelectron microscopic analysis and proteolytic digestion, to be in the outer membrane fraction as well as in the cytoplasm (173), surface exposed (170, 172), and in outer membrane vesicles (OMVs) shed by *B. burgdorferi* during culture (171). The enolase in the OMV is accessible to proteolytic degradation by proteinase K, providing evidence for its surface exposure in OMVs. Moreover, the importance of enolase in infection is implied by its higher production under conditions that mimic the conditions in the mammalian host (171). Taken together, these findings suggest a moonlighting role for enolase in *B. burgdorferi* by virtue of its role in stabilizing cell-bound or extracellular OMV-bound plasmin in addition to its canonical role in glycolysis. The binding of PLG to bacterial enolases is not restricted to *Borrelia*. *Streptococcus canis* enolase binds to both human and canine plasminogen and facilitates degradation of fibrin after activation with host-derived uPA (174). *Leptospira interrogans* binds PLG in what appears to be a unique manner, whereby the enolase is secreted, binds PLG, and reassociates with the organism (172).

Other *B. burgdorferi* outer surface proteins have been identified as interacting with PLG. Assays using radiolabeled human PLG to bind *B. burgdorferi* identified two prominent 70- and 30-kDa PLG-binding outer membrane proteins, of which, the 30-kDa band was found to be OspA (152). The 70-kDa band, termed BPBP, adsorbed approximately 10 times more label than OspA (175). An interesting and unexpected finding was that there was seemingly no difference between low-passage-number (infectious) and high-passage-number strains in the binding of PLG (152, 175). Genetic alignments showed BPBP to be an ortholog of a 60-kDa protein found in *B. coriaceae* (66% identity, 80% similarity). This protein was antigenic in infected patients (175). In another study, recombinant hypothetical protein BBA70 bound PLG in a dose-dependent manner, and the bound plasmin degraded chromogenic substrate S-2251 and natural substrate fibrinogen as well as complement system components C3b and C5 (150).

Interaction of *Borrelia* with Other PAS Components

As mentioned earlier, *Borrelia* spp. interact with each of the components of the PAS (Fig. 3). *Borrelia* spp. interact with the matrix metalloprotease system (Fig. 3) by inducing the production of these proteases from a diverse group of cells. The level of interaction of *Borrelia* and all the components of the PAS is thorough and meets all of its physiological requirements. Lyme and relapsing fever *Borrelia* spp. stimulated human peripheral blood monocytes and neutrophils to release pro-matrix metalloproteinase-9 (pro-MMP-9; gelatinase B) and active matrix metalloproteinase-1 (MMP-1; collagenase-1). U937 cells and human keratinocyte cells released pro-MMP-9, and plasmin stabilized on the surface of the Lyme disease spirochete activated pro-MMP-9 to its active form. This active form was also observed in the plasma of mice infected with a relapsing fever *Borrelia*. These results showed that *Borrelia* can upregulate MMPs and mediate an activation cascade initiated by plasmin bound to the microbial surface. *B. burgdorferi*-bound plasmin triggered the conversion of pro-matrix metalloproteases (MMPs) to active MMP-9 (type IV collagenase), establishing an indirect role for plasmin in collagen degradation (176).

Attempts at working out the mechanisms for production of MMPs by the spirochetes is tied to proinflammatory pathways. Secretion of MMP-9 by *B. burgdorferi* is selectively induced through Toll-like receptor (TLR) 2 in human and murine monocytic cells (176). Secretion of MMP-1 was shown to be stimulated through a pathway other than TLR2 (177). *B. burgdorferi* utilizes several signal transduction pathways for the production of MMPs from human joint cells (178). *B. burgdorferi* induced transcription of MMP-1, -3, -13, and -19 from primary human chondrocyte cells (178). Transcription of MMP-10 and tissue inhibitor of MMP-1 were also elevated but without concurrent protein expression. The synovial fluid levels of MMPs in patients matched the experimental results. In contrast, infection of mice with *B. burgdorferi* only induced production of

MMP-3 and MMP-19 (179). *B. burgdorferi* binds to integrin $\alpha(3)\beta(1)$, and binding of this complex induces production of matrix metalloproteinases in primary human chondrocyte cells (180).

MMPs have also been detected in clinical specimens, emphasizing their role in Lyme disease pathogenesis. Not surprisingly, MMPs have been associated with the skin, neurological, and joint manifestations. MMP-9 was detected in the cerebrospinal fluid (CSF) of patients with neuroborreliosis, and infiltrating macrophages were thought to be a possible source of the MMP-9 increase (181). Levels of MMP-1 and MMP-3 in synovial fluid were higher in untreated Lyme arthritis patients. In contrast, MMP-8 and MMP-9 were elevated in patients with antibiotic-resistant arthritis, illustrating the diversity of stimulation pathways depending on disease stage (182). MMP-9 was upregulated in erythema migrans lesions (183). Possible mechanisms for MMP induction have been proposed as well. Levels of MMP-9 and soluble CD14 were markedly elevated in serum from patients and were also upregulated in U937 cells by *B. burgdorferi* in a time- and concentration-dependent manner. When fibroblasts incubated with supernatants from U937 cells were exposed to *B. burgdorferi* or recombinant CD14, the expression of MMP-9 was significantly increased (184).

As previously stated, the *Borrelia* spp. interact with all the components of the PAS (Fig. 3). In coculture experiments with monocytic cells, *B. burgdorferi*, as well as various spirochetal components, induced the expression of cellular uPA receptor (uPAR) as well as the release of a soluble form. The upregulation of uPAR by *B. burgdorferi* was concomitant with an increase in uPAR mRNA. The mechanism for soluble uPAR release and its role were not apparent in this study (185). In an ensuing study with monocytic cells, combined functional blockade of CD14 and TLR2 results in significant, but not total, inhibition of *B. burgdorferi*-dependent uPAR induction. It was concluded that signaling for monocytic cell uPAR expression mediated by *B. burgdorferi* proceeds with CD14 and TLR2 as partial contributors. That part, under the control of CD14 and TLR2, represents a link between the host PLG activation and innate immunity systems (186).

Borrowed Proteolysis and Host Defense

The most accepted (perhaps the only) outcome of having active plasmin on the surface of *Borrelia* is increased invasiveness. Resulting proteolysis of matrix proteins and connective tissue by bound plasmin can assist this highly motile organism in invading tissues. Likewise, proteolysis of the external milieu of the spirochete by bound plasmin could be an important means for nutrient acquisition, and this is an area that merits future investigation.

What if PLG binding followed by uPA activation was a host defense mechanism against the bacteria that utilize the PAS? There are scant but tantalizing indications that this could be the case. One such early indication comes from the stimulation of uPAR production by *B. burgdorferi* (185, 186). In a far-reaching study, uPAR knockout mice had significantly higher numbers of *B. burgdorferi* than the controls (187). These mice also displayed impaired phagocytic capacity. In uPAR knockout mice backcrossed to a C3H/HeN background (the preferred inbred mouse strain for laboratory infection), the higher numbers of spirochetes were associated with more severe carditis and increased local TLR2 and interleukin 1 beta (IL-1 β) expression. This study concluded that in *B. burgdorferi* infection, uPAR is required for phagocytosis and adequate eradication of the spirochete from the heart by a mechanism that is independent of the binding of uPAR to uPA or its role in the fibrinolytic system (187, 188). Moreover, the recently discovered PLG receptor on the surfaces of cells of the innate immune system (189) is important in the PLG-dependent regulation of the migration, invasion, and recruitment of macrophages and inflammatory monocytes (190, 191). This is an expanding area of investigation (188) that could have an impact on the host defense against PLG-bound bacteria.

CONCLUDING REMARKS

In this review, we have considered the protease contingent of *Borrelia*, from annotation in the genome to the research on specific proteases of these spirochetes. The




FIG 4 A schematic overview of the key endogenous and borrowed proteases *Borrelia* spp. use to remodel their proteomes as they move from one stage, tissue, or host to another. Major cytoplasmic and inner membrane proteases, Lon, ClpAP (as a representative of the Clp protease systems ClpXP, ClpYQ, and ClpCP), and FtsH are shown. There are bound substrates highlighted in red. The lone periplasmic protease HtrA, which is also secreted, is shown in the periplasmic space and associated with the cell surface. The yellow ovals represent cholesterol-rich lipid rafts, which are of functional importance and exist in both the inner and outer membranes of *Borrelia*. The borrowed plasmin proteolytic system is shown bound to *Borrelia* outer surface proteins (Osp), and additional cell surface receptors, shown as a green oval. The 5 kringle domains of plasmin are colored purple, and its protease domain is light green. The periplasmic flagella are not shown.

review has shown some unique features of the proteolytic arsenal (Fig. 4) and the realization that *Borrelia* spp. (both RF and Lyme species) have dedicated a large percentage of their genomes to proteolysis. The proteolytic activities of secreted HtrA have led to new paradigms of pathogenesis, and we reviewed this aspect in the context of other bacteria. The interactions of *Borrelia* with all the components of the plasminogen activation system is an example of pathogens appropriating host proteolytic activities. Borrowed proteolysis and its role in pathogenesis, not only by infection with *Borrelia* but also by infection with other bacteria, could be viewed as a mechanism of host defense. One of the major reasons for writing this review was to consider the overwhelming evidence of the utilization of the PAS by *Borrelia*. There is no other mammalian system so intertwined with a group of pathogens such as the RF and LD *Borrelia* spp. Moreover, the evidence for the utilization of the PAS stems from experimental *in vitro* studies to *in vivo* mouse model experiments and to actual studies with patients with borrelioses. Despite this, there is much that needs to be done, as surprisingly little is known about the targets of the spirochetal harbored proteases, their temporal regulation, and their unique mechanisms of action. Likewise, a reassessment of borrowed host proteolysis could reveal surprising consequences for pathogenesis.

ACKNOWLEDGMENTS

J.L.B. and A.W.K. are supported by the U.S. National Institutes of Health through extramural research grants AI-027044 and AI-127533, respectively.

There are no financial interests or support from institutions or companies mentioned in the manuscript.

J.L.C., J.L.B., and A.W.K. researched data for the article, substantially contributed to discussion of the content, wrote, reviewed, and edited the manuscript.

REFERENCES

- Swelengrebel NH. 1907. De l'anaphylaxie en général. *Ann Inst Pasteur* 21:562–586.
- Cutler SJ, Ruzic-Sabljic E, Potkonjak A. 2017. Emerging borreliae - expanding beyond Lyme borreliosis. *Mol Cell Probes* 31:22–27. <https://doi.org/10.1016/j.mcp.2016.08.003>.
- Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. 1982. Lyme disease—a tick-borne spirochetosis? *Science* 216:1317–1319. <https://doi.org/10.1126/science.7043737>.
- Fukunaga M, Takahashi Y, Tsuruta Y, Matsushita O, Ralph D, McClelland M, Nakao M. 1995. Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov., isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int J Syst Bacteriol* 45:804–810. <https://doi.org/10.1099/00207713-45-4-804>.
- Charon NW, Cockburn A, Li C, Liu J, Miller KA, Miller MR, Motaleb MA, Wolgemuth CW. 2012. The unique paradigm of spirochete motility and chemotaxis. *Annu Rev Microbiol* 66:349–370. <https://doi.org/10.1146/annurev-micro-092611-150145>.
- Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb JF, Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J, Weidman J, Utterback T, Watthey L, McDonald L, Artiach P, Bowman C, Garland S, Fuji C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter JC. 1997. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. *Nature* 390:580–586. <https://doi.org/10.1038/37551>.
- LaRocca TJ, Crowley JT, Cusack BJ, Pathak P, Benach J, London E, Garcia-Monco JC, Benach JL. 2010. Cholesterol lipids of *Borrelia burgdorferi* form lipid rafts and are required for the bactericidal activity of a complement-independent antibody. *Cell Host Microbe* 8:331–342. <https://doi.org/10.1016/j.chom.2010.09.001>.
- Radolf JD, Samuels DS. 2021. Lyme disease and relapsing fever spirochetes. Caister Academic Press, Poole, United Kingdom. <https://doi.org/10.21775/9781913652616>.
- Radolf JD, Caimano MJ, Stevenson B, Hu LT. 2012. Of ticks, mice and men: understanding the dual-host lifestyle of Lyme disease spirochaetes. *Nat Rev Microbiol* 10:87–99. <https://doi.org/10.1038/nrmicro2714>.
- Benach JL, Bosler EM, Hanrahan JP, Coleman JL, Habicht GS, Bast TF, Cameron DJ, Ziegler JL, Barbour AG, Burgdorfer W, Edelman R, Kaslow RA. 1983. Spirochetes isolated from the blood of two patients with Lyme disease. *N Engl J Med* 308:740–742. <https://doi.org/10.1056/NEJM198303313081302>.
- Steere AC, Grodzicki RL, Kornblatt AN, Craft JE, Barbour AG, Burgdorfer W, Schmid GP, Johnson E, Malawista SE. 1983. The spirochetal etiology of Lyme disease. *N Engl J Med* 308:733–740. <https://doi.org/10.1056/NEJM198303313081301>.
- Hedstrom L. 2002. Serine protease mechanism and specificity. *Chem Rev* 102:4501–4524. <https://doi.org/10.1021/cr000033x>.
- Hooper NM. 2002. Proteases: a primer. *Essays Biochem* 38:1–8. <https://doi.org/10.1042/bse0380001>.
- Klein T, Eckhard U, Dufour A, Solis N, Overall CM. 2018. Proteolytic cleavage-mechanisms, function, and “omic” approaches for a near-ubiquitous posttranslational modification. *Chem Rev* 118:1137–1168. <https://doi.org/10.1021/acs.chemrev.7b00120>.
- Lopez-Otin C, Bond JS. 2008. Proteases: multifunctional enzymes in life and disease. *J Biol Chem* 283:30433–30437. <https://doi.org/10.1074/jbc.R800035200>.
- Paetzel M. 2019. Bacterial signal peptidases. *Subcell Biochem* 92:187–219. https://doi.org/10.1007/978-3-030-18768-2_7.
- Grab DJ, Kennedy R, Philipp MT. 1996. *Borrelia burgdorferi* possesses a collagenolytic activity. *FEMS Microbiol Lett* 144:39–45. <https://doi.org/10.1111/j.1574-6968.1996.tb08506.x>.
- Liu J, Mei Z, Li N, Qi Y, Xu Y, Shi Y, Wang F, Lei J, Gao N. 2013. Structural dynamics of the MecA-ClpC complex: a type II AAA+ protein unfolding machine. *J Biol Chem* 288:17597–17608. <https://doi.org/10.1074/jbc.M113.458752>.
- Stahlhut SG, Alqarzaee AA, Jensen C, Fisker NS, Pereira AR, Pinho MG, Thomas VC, Frees D. 2017. The ClpXP protease is dispensable for degradation of unfolded proteins in *Staphylococcus aureus*. *Sci Rep* 7:11739. <https://doi.org/10.1038/s41598-017-12122-y>.
- Kebe NM, Samanta K, Singh P, Lai-Kee-Him J, Apicella V, Payrot N, Lauraire N, Legrand B, Lisowski V, Mbang-Benet DE, Pages M, Bastien P, Kajava AV, Bron P, Hernandez JF, Coux O. 2019. The HslV protease from *Leishmania major* and its activation by C-terminal HslU peptides. *Int J Mol Sci* 20:1021. <https://doi.org/10.3390/ijms20051021>.
- Sundar S, Baker TA, Sauer RT. 2012. The I domain of the AAA+ HslUV protease coordinates substrate binding, ATP hydrolysis, and protein degradation. *Protein Sci* 21:188–198. <https://doi.org/10.1002/pro.2001>.
- Ogden AJ, McAleer JM, Kahn ML. 2019. Characterization of the *Sinorhizobium meliloti* HslUV and ClpXP protease systems in free-living and symbiotic states. *J Bacteriol* 201:e00498-18. <https://doi.org/10.1128/JB.00498-18>.
- Bertin PB, Lozzi SP, Howell JK, Restrepo-Cadavid G, Neves D, Teixeira AR, de Sousa MV, Norris SJ, Santana JM. 2005. The thermophilic, homohexameric aminopeptidase of *Borrelia burgdorferi* is a member of the M29 family of metalloproteases. *Infect Immun* 73:2253–2261. <https://doi.org/10.1128/IAI.73.4.2253-2261.2005>.
- Ostberg Y, Carroll JA, Pinne M, Krum JG, Rosa P, Bergstrom S. 2004. Pleiotropic effects of inactivating a carboxyl-terminal protease, CtpA, in *Borrelia burgdorferi*. *J Bacteriol* 186:2074–2084. <https://doi.org/10.1128/jb.186.7.2074-2084.2004>.
- Kumru OS, Bunikis I, Sorokina I, Bergstrom S, Zuckert WR. 2011. Specificity and role of the *Borrelia burgdorferi* CtpA protease in outer membrane protein processing. *J Bacteriol* 193:5759–5765. <https://doi.org/10.1128/JB.05622-11>.
- Chu CY, Stewart PE, Bestor A, Hansen B, Lin T, Gao L, Norris SJ, Rosa PA. 2016. Function of the *Borrelia burgdorferi* FtsH homolog is essential for viability both *in vitro* and *in vivo* and independent of HflK/C. *mBio* 7:e00404-16. <https://doi.org/10.1128/mBio.00404-16>.
- van Stelten J, Silva F, Belin D, Silhavy TJ. 2009. Effects of antibiotics and a proto-oncogene homolog on destruction of protein translocator SecY. *Science* 325:753–756. <https://doi.org/10.1126/science.1172221>.
- Krzywda S, Brzozowski AM, Verma C, Karata K, Ogura T, Wilkinson AJ. 2002. The crystal structure of the AAA domain of the ATP-dependent protease FtsH of *Escherichia coli* at 1.5 Å resolution. *Structure* 10:1073–1083. [https://doi.org/10.1016/s0969-2126\(02\)00806-7](https://doi.org/10.1016/s0969-2126(02)00806-7).
- Schumann W. 1999. FtsH—a single-chain charonin? *FEMS Microbiol Rev* 23:1–11. <https://doi.org/10.1111/j.1574-6976.1999.tb00389.x>.
- Santos D, De Almeida DF. 1975. Isolation and characterization of a new temperature-sensitive cell division mutant of *Escherichia coli* K-12. *J Bacteriol* 124:1502–1507. <https://doi.org/10.1128/JB.124.3.1502-1507.1975>.
- Saikawa N, Akiyama Y, Ito K. 2004. FtsH exists as an exceptionally large complex containing HflK in the plasma membrane of *Escherichia coli*. *J Struct Biol* 146:123–129. <https://doi.org/10.1016/j.jsb.2003.09.020>.
- Toledo A, Huang Z, Coleman JL, London E, Benach JL. 2018. Lipid rafts can form in the inner and outer membranes of *Borrelia burgdorferi* and have different properties and associated proteins. *Mol Microbiol* 108:63–76. <https://doi.org/10.1111/mmi.13914>.
- Toledo A, Crowley JT, Coleman JL, LaRocca TJ, Chiantia S, London E, Benach JL. 2014. Selective association of outer surface lipoproteins with the lipid rafts of *Borrelia burgdorferi*. *mBio* 5:e00899-14. <https://doi.org/10.1128/mBio.00899-14>.
- He L, Luo D, Yang F, Li C, Zhang X, Deng H, Zhang JR. 2018. Multiple domains of bacterial and human Lon proteases define substrate selectivity. *Emerg Microbes Infect* 7:149. <https://doi.org/10.1038/s41426-018-0148-4>.
- Liao JH, Lin YC, Hsu J, Lee AY, Chen TA, Hsu CH, Chir JL, Hua KF, Wu TH, Hong LJ, Yen PW, Chiou A, Wu SH. 2010. Binding and cleavage of *E. coli* HUBeta by the *E. coli* Lon protease. *Biophys J* 98:129–137. <https://doi.org/10.1016/j.bpj.2009.09.052>.
- Van Melder L, Thi MH, Lecchi P, Gottesman S, Couturier M, Maurizi MR. 1996. ATP-dependent degradation of CcdA by Lon protease. Effects of secondary structure and heterologous subunit interactions. *J Biol Chem* 271:27730–27738. <https://doi.org/10.1074/jbc.271.44.27730>.
- Bissonnette SA, Rivera-Rivera I, Sauer RT, Baker TA. 2010. The IbpA and IbpB small heat-shock proteins are substrates of the AAA+ Lon protease. *Mol Microbiol* 75:1539–1549. <https://doi.org/10.1111/j.1365-2958.2010.07070.x>.
- Shah IM, Wolf RE, Jr. 2006. Sequence requirements for Lon-dependent degradation of the *Escherichia coli* transcription activator SoxS: identification of the SoxS residues critical to proteolysis and specific inhibition of *in vitro* degradation by a peptide comprised of the N-terminal 21 amino acid residues. *J Mol Biol* 357:718–731. <https://doi.org/10.1016/j.jmb.2005.12.088>.
- Mukherjee S, Bree AC, Liu J, Patrick JE, Chien P, Kearns DB. 2015. Adaptor-mediated Lon proteolysis restricts *Bacillus subtilis* hyperflagellation.

- Proc Natl Acad Sci U S A 112:250–255. <https://doi.org/10.1073/pnas.1417419112>.
40. Choy JS, Aung LL, Karzai AW. 2007. Lon protease degrades transfer-messenger RNA-tagged proteins. *J Bacteriol* 189:6564–6571. <https://doi.org/10.1128/JB.00860-07>.
 41. Karzai AW, Roche ED, Sauer RT. 2000. The SsrA-SmpB system for protein tagging, directed degradation and ribosome rescue. *Nat Struct Biol* 7:449–455. <https://doi.org/10.1038/75843>.
 42. Puri N, Karzai AW. 2017. HspQ Functions as a unique specificity-enhancing factor for the AAA+ Lon protease. *Mol Cell* 66:672.e4–683.e4. <https://doi.org/10.1016/j.molcel.2017.05.016>.
 43. Choi J, Groisman EA. 2020. *Salmonella* expresses foreign genes during infection by degrading their silencer. *Proc Natl Acad Sci U S A* 117:8074–8082. <https://doi.org/10.1073/pnas.1912808117>.
 44. Cloud JL, Marconi RT, Eggers CH, Garon CF, Tilly K, Samuels DS. 1997. Cloning and expression of the *Borrelia burgdorferi* lon gene. *Gene* 194:137–141. [https://doi.org/10.1016/S0378-1119\(97\)00196-0](https://doi.org/10.1016/S0378-1119(97)00196-0).
 45. Tokarz R, Anderton JM, Katona LI, Benach JL. 2004. Combined effects of blood and temperature shift on *Borrelia burgdorferi* gene expression as determined by whole genome DNA array. *Infect Immun* 72:5419–5432. <https://doi.org/10.1128/IAI.72.9.5419-5432.2004>.
 46. Coleman JL, Katona LI, Kuhlow C, Toledo A, Okan NA, Tokarz R, Benach JL. 2009. Evidence that two ATP-dependent (Lon) proteases in *Borrelia burgdorferi* serve different functions. *PLoS Pathog* 5:e1000676. <https://doi.org/10.1371/journal.ppat.1000676>.
 47. Thompson C, Mason C, Parrilla S, Ouyang Z. 2020. The Lon-1 protease is required by *Borrelia burgdorferi* to infect the mammalian host. *Infect Immun* 88:e00951-19. <https://doi.org/10.1128/IAI.00951-19>.
 48. Mason C, Thompson C, Ouyang Z. 2020. The Lon-2 protease of *Borrelia burgdorferi* is critical for infection in the mammalian host. *Mol Microbiol* 113:938–950. <https://doi.org/10.1111/mmi.14460>.
 49. Clausen T, Kaiser M, Huber R, Ehrmann M. 2011. HTRA proteases: regulated proteolysis in protein quality control. *Nat Rev Mol Cell Biol* 12:152–162. <https://doi.org/10.1038/nrm3065>.
 50. Swamy KH, Chung CH, Goldberg AL. 1983. Isolation and characterization of protease do from *Escherichia coli*, a large serine protease containing multiple subunits. *Arch Biochem Biophys* 224:543–554. [https://doi.org/10.1016/0003-9861\(83\)90242-4](https://doi.org/10.1016/0003-9861(83)90242-4).
 51. Ehrmann M, Clausen T. 2004. Proteolysis as a regulatory mechanism. *Annu Rev Genet* 38:709–724. <https://doi.org/10.1146/annurev.genet.38.072902.093416>.
 52. Merdanovic M, Clausen T, Kaiser M, Huber R, Ehrmann M. 2011. Protein quality control in the bacterial periplasm. *Annu Rev Microbiol* 65:149–168. <https://doi.org/10.1146/annurev-micro-090110-102925>.
 53. Krojer T, Garrido-Franco M, Huber R, Ehrmann M, Clausen T. 2002. Crystal structure of DegP (HtrA) reveals a new protease-chaperone machine. *Nature* 416:455–459. <https://doi.org/10.1038/416455a>.
 54. Krojer T, Sawa J, Schafer E, Saibil HR, Ehrmann M, Clausen T. 2008. Structural basis for the regulated protease and chaperone function of DegP. *Nature* 453:885–890. <https://doi.org/10.1038/nature07004>.
 55. Guyard C, Battisti JM, Raffel SJ, Schruppf ME, Whitney AR, Krum JG, Porcella SF, Rosa PA, DeLeo FR, Schwan TG. 2006. Relapsing fever spirochaetes produce a serine protease that provides resistance to oxidative stress and killing by neutrophils. *Mol Microbiol* 60:710–722. <https://doi.org/10.1111/j.1365-2958.2006.05122.x>.
 56. Jackson-Litteken CD, Zalud AK, Ratliff CT, Latham JI, Bourret TJ, Lopez JE, Blevins JS. 2019. Assessing the contribution of an HtrA family serine protease during *Borrelia turicatae* mammalian infection. *Front Cell Infect Microbiol* 9:290. <https://doi.org/10.3389/fcimb.2019.00290>.
 57. Kariu T, Yang X, Marks CB, Zhang X, Pal U. 2013. Proteolysis of BB0323 results in two polypeptides that impact physiologic and infectious phenotypes in *Borrelia burgdorferi*. *Mol Microbiol* 88:510–522. <https://doi.org/10.1111/mmi.12202>.
 58. Zhang X, Yang X, Kumar M, Pal U. 2009. BB0323 function is essential for *Borrelia burgdorferi* virulence and persistence through tick-rodent transmission cycle. *J Infect Dis* 200:1318–1330. <https://doi.org/10.1086/605846>.
 59. Coleman JL, Crowley JT, Toledo AM, Benach JL. 2013. The HtrA protease of *Borrelia burgdorferi* degrades outer membrane protein BmpD and chemotaxis phosphatase CheX. *Mol Microbiol* 88:619–633. <https://doi.org/10.1111/mmi.12213>.
 60. Russell TM, Tang X, Goldstein JM, Bagarozzi D, Johnson BJ. 2016. The salt-sensitive structure and zinc inhibition of *Borrelia burgdorferi* protease BbHtrA. *Mol Microbiol* 99:586–596. <https://doi.org/10.1111/mmi.13251>.
 61. Zhang K, Qin Z, Chang Y, Liu J, Malkowski MG, Shipa S, Li L, Qiu W, Zhang JR, Li C. 2019. Analysis of a flagellar filament cap mutant reveals that HtrA serine protease degrades unfolded flagellin protein in the periplasm of *Borrelia burgdorferi*. *Mol Microbiol* 111:1652–1670. <https://doi.org/10.1111/mmi.14243>.
 62. Toledo A, Perez A, Coleman JL, Benach JL. 2015. The lipid raft proteome of *Borrelia burgdorferi*. *Proteomics* 15:3662–3675. <https://doi.org/10.1002/pmic.201500093>.
 63. Coleman JL, Toledo A, Benach JL. 2016. *Borrelia burgdorferi* HtrA: evidence for twofold proteolysis of outer membrane protein p66. *Mol Microbiol* 99:135–150. <https://doi.org/10.1111/mmi.13221>.
 64. Kobryn K, Naigamwalla DZ, Chaconas G. 2000. Site-specific DNA binding and bending by the *Borrelia burgdorferi* Hbb protein. *Mol Microbiol* 37:145–155. <https://doi.org/10.1046/j.1365-2958.2000.01981.x>.
 65. Coburn J, Cugini C. 2003. Targeted mutation of the outer membrane protein P66 disrupts attachment of the Lyme disease agent, *Borrelia burgdorferi*, to integrin alphavbeta3. *Proc Natl Acad Sci U S A* 100:7301–7306. <https://doi.org/10.1073/pnas.113117100>.
 66. Barcena-Uribarri I, Thein M, Sacher A, Bunikis I, Bonde M, Bergstrom S, Benz R. 2010. P66 porins are present in both Lyme disease and relapsing fever spirochetes: a comparison of the biophysical properties of P66 porins from six *Borrelia* species. *Biochim Biophys Acta* 1798:1197–1203. <https://doi.org/10.1016/j.bbame.2010.02.011>.
 67. Kenedy MR, Luthra A, Anand A, Dunn JP, Radolf JD, Akins DR. 2014. Structural modeling and physicochemical characterization provide evidence that P66 forms a beta-barrel in the *Borrelia burgdorferi* outer membrane. *J Bacteriol* 196:859–872. <https://doi.org/10.1128/JB.01236-13>.
 68. Coleman JL, Toledo A, Benach JL. 2018. HtrA of *Borrelia burgdorferi* leads to decreased swarm motility and decreased production of pyruvate. *mBio* 9:e01136-18. <https://doi.org/10.1128/mBio.01136-18>.
 69. Zhuang X, Yang X, Altieri AS, Nelson DC, Pal U. 2018. *Borrelia burgdorferi* surface-located Lmp1 protein processed into region-specific polypeptides that are critical for microbial persistence. *Cell Microbiol* 20:e12855. <https://doi.org/10.1111/cmi.12855>.
 70. Russell TM, Delorey MJ, Johnson BJ. 2013. *Borrelia burgdorferi* BbHtrA degrades host ECM proteins and stimulates release of inflammatory cytokines *in vitro*. *Mol Microbiol* 90:241–251. <https://doi.org/10.1111/mmi.12377>.
 71. Russell TM, Johnson BJ. 2013. Lyme disease spirochaetes possess an aggrecan-binding protease with aggrecanase activity. *Mol Microbiol* 90:228–240. <https://doi.org/10.1111/mmi.12276>.
 72. Ullmann AJ, Russell TM, Dolan MC, Williams M, Hojgaard A, Weiner ZP, Johnson BJ. 2015. Evaluation of *Borrelia burgdorferi* BbHtrA protease as a vaccine candidate for Lyme borreliosis in mice. *PLoS One* 10:e0128868. <https://doi.org/10.1371/journal.pone.0128868>.
 73. Ye M, Sharma K, Thakur M, Smith AA, Buyuktanir O, Xiang X, Yang X, Promnares K, Lou Y, Yang XF, Pal U. 2016. HtrA, a temperature- and stationary phase-activated protease involved in maturation of a key microbial virulence determinant, facilitates *Borrelia burgdorferi* infection in mammalian hosts. *Infect Immun* 84:2372–2381. <https://doi.org/10.1128/IAI.00360-16>.
 74. Boehm M, Simson D, Escher U, Schmidt AM, Bereswill S, Tegtmeyer N, Backert S, Heimesaat MM. 2018. Function of serine protease HtrA in the lifecycle of the foodborne pathogen *Campylobacter jejuni*. *Eur J Microbiol Immunol (Bp)* 8:70–77. <https://doi.org/10.1556/1886.2018.00011>.
 75. Hoy B, Lower M, Weydig C, Carra G, Tegtmeyer N, Geppert T, Schroder P, Sewald N, Backert S, Schneider G, Wessler S. 2010. *Helicobacter pylori* HtrA is a new secreted virulence factor that cleaves E-cadherin to disrupt intercellular adhesion. *EMBO Rep* 11:798–804. <https://doi.org/10.1038/embor.2010.114>.
 76. Zarzecka U, Modrak-Wojcik A, Figaj D, Apanowicz M, Lesner A, Bzowska A, Lipinska B, Zawilak-Pawlik A, Backert S, Skorko-Glonek J. 2019. Properties of the HtrA protease from bacterium *Helicobacter pylori* whose activity is indispensable for growth under stress conditions. *Front Microbiol* 10:961. <https://doi.org/10.3389/fmicb.2019.00961>.
 77. Backert S, Bernegger S, Skorko-Glonek J, Wessler S. 2018. Extracellular HtrA serine proteases: an emerging new strategy in bacterial pathogenesis. *Cell Microbiol* 20:e12845. <https://doi.org/10.1111/cmi.12845>.

78. Wiman B. 1973. Primary structure of peptides released during activation of human plasminogen by urokinase. *Eur J Biochem* 39:1–9. <https://doi.org/10.1111/j.1432-1033.1973.tb03096.x>.
79. Lijnen HR. 2001. Elements of the fibrinolytic system. *Ann N Y Acad Sci* 936:226–236. <https://doi.org/10.1111/j.1749-6632.2001.tb03511.x>.
80. Dano K, Andreasen PA, Grondahl-Hansen J, Kristensen P, Nielsen LS, Skriver L. 1985. Plasminogen activators, tissue degradation, and cancer. *Adv Cancer Res* 44:139–266. [https://doi.org/10.1016/s0065-230x\(08\)60028-7](https://doi.org/10.1016/s0065-230x(08)60028-7).
81. Pedersen N, Schmitt M, Ronne E, Nicoletti MI, Hoyer-Hansen G, Conese M, Giavazzi R, Dano K, Kuhn W, Janicke F. 1993. A ligand-free, soluble urokinase receptor is present in the ascitic fluid from patients with ovarian cancer. *J Clin Invest* 92:2160–2167. <https://doi.org/10.1172/JCI116817>.
82. Fazioli F, Blasi F. 1994. Urokinase-type plasminogen activator and its receptor: new targets for anti-metastatic therapy? *Trends Pharmacol Sci* 15:25–29. [https://doi.org/10.1016/0165-6147\(94\)90130-9](https://doi.org/10.1016/0165-6147(94)90130-9).
83. Tang L, Han X. 2013. The urokinase plasminogen activator system in breast cancer invasion and metastasis. *Biomed Pharmacother* 67:179–182. <https://doi.org/10.1016/j.biopha.2012.10.003>.
84. Plow EF, Freaney DE, Plescia J, Miles LA. 1986. The plasminogen system and cell surfaces: evidence for plasminogen and urokinase receptors on the same cell type. *J Cell Biol* 103:2411–2420. <https://doi.org/10.1083/jcb.103.6.2411>.
85. Lamba D, Bauer M, Huber R, Fischer S, Rudolph R, Kohnert U, Bode W. 1996. The 2.3 Å crystal structure of the catalytic domain of recombinant two-chain human tissue-type plasminogen activator. *J Mol Biol* 258:117–135. <https://doi.org/10.1006/jmbi.1996.0238>.
86. Vassalli JD, Baccino D, Belin D. 1985. A cellular binding site for the Mr 55,000 form of the human plasminogen activator, urokinase. *J Cell Biol* 100:86–92. <https://doi.org/10.1083/jcb.100.1.86>.
87. Stoppelli MP, Corti A, Soffientini A, Cassani G, Blasi F, Assoian RK. 1985. Differentiation-enhanced binding of the amino-terminal fragment of human urokinase plasminogen activator to a specific receptor on U937 monocytes. *Proc Natl Acad Sci U S A* 82:4939–4943. <https://doi.org/10.1073/pnas.82.15.4939>.
88. Blasi F, Stoppelli MP, Cubellis MV. 1986. The receptor for urokinase-plasminogen activator. *J Cell Biochem* 32:179–186. <https://doi.org/10.1002/jcb.240320303>.
89. Min HY, Semnani R, Mizukami IF, Watt K, Todd RF, III, Liu DY. 1992. cDNA for Mo3, a monocyte activation antigen, encodes the human receptor for urokinase plasminogen activator. *J Immunol* 148:3636–3642.
90. Cubellis MV, Noll ML, Cassani G, Blasi F. 1986. Binding of single-chain prourokinase to the urokinase receptor of human U937 cells. *J Biol Chem* 261:15819–15822. [https://doi.org/10.1016/S0021-9258\(18\)66635-9](https://doi.org/10.1016/S0021-9258(18)66635-9).
91. Nielsen LS, Kellerman GM, Behrendt N, Picone R, Dano K, Blasi F. 1988. A 55,000–60,000 Mr receptor protein for urokinase-type plasminogen activator. Identification in human tumor cell lines and partial purification. *J Biol Chem* 263:2358–2363. [https://doi.org/10.1016/S0021-9258\(18\)69214-2](https://doi.org/10.1016/S0021-9258(18)69214-2).
92. Wei Y, Lukashev M, Simon DI, Bodary SC, Rosenberg S, Doyle MV, Chapman HA. 1996. Regulation of integrin function by the urokinase receptor. *Science* 273:1551–1555. <https://doi.org/10.1126/science.273.5281.1551>.
93. Mahmood N, Mihalicioiu C, Rabbani SA. 2018. Multifaceted role of the urokinase-type plasminogen activator (uPA) and its receptor (uPAR): diagnostic, prognostic, and therapeutic applications. *Front Oncol* 8:24. <https://doi.org/10.3389/fonc.2018.00024>.
94. Castellino FJ, Ploplis VA. 2005. Structure and function of the plasminogen/plasmin system. *Thromb Haemost* 93:647–654. <https://doi.org/10.1160/TH04-12-0842>.
95. Law RH, Abu-Ssaydeh D, Whistock JC. 2013. New insights into the structure and function of the plasminogen/plasmin system. *Curr Opin Struct Biol* 23:836–841. <https://doi.org/10.1016/j.sbi.2013.10.006>.
96. Redlitz A, Plow EF. 1995. Receptors for plasminogen and t-PA: an update. *Baillieres Clin Haematol* 8:313–327. [https://doi.org/10.1016/s0950-3536\(05\)80270-7](https://doi.org/10.1016/s0950-3536(05)80270-7).
97. Collen D, Lijnen HR. 2005. Thrombolytic agents. *Thromb Haemost* 93:627–630. <https://doi.org/10.1160/TH04-11-0724>.
98. Lee AY, Fredenburgh JC, Stewart RJ, Rischke JA, Weitz JI. 2001. Like fibrin, (DD)E, the major degradation product of crosslinked fibrin, protects plasmin from inhibition by alpha2-antiplasmin. *Thromb Haemost* 85:502–508. <https://doi.org/10.1055/s-0037-1615612>.
99. Schneider M, Nesheim M. 2004. A study of the protection of plasmin from antiplasmin inhibition within an intact fibrin clot during the course of clot lysis. *J Biol Chem* 279:13333–13339. <https://doi.org/10.1074/jbc.M313164200>.
100. Travis J, Salvesen GS. 1983. Human plasma proteinase inhibitors. *Annu Rev Biochem* 52:655–709. <https://doi.org/10.1146/annurev.bi.52.070183.003255>.
101. Zwirzitz A, Reiter M, Skrabana R, Ohradanova-Repic A, Majdic O, Gutekova M, Cehlar O, Petrovcikova E, Kutejova E, Stanek G, Stockinger H, Leksa V. 2018. Lactoferrin is a natural inhibitor of plasminogen activation. *J Biol Chem* 293:8600–8613. <https://doi.org/10.1074/jbc.RA118.003145>.
102. Reddy KN, Markus G. 1972. Mechanism of activation of human plasminogen by streptokinase. Presence of active center in streptokinase-plasminogen complex. *J Biol Chem* 247:1683–1691. [https://doi.org/10.1016/S0021-9258\(19\)45531-2](https://doi.org/10.1016/S0021-9258(19)45531-2).
103. Castellino FJ, Sodetz JM, Brockway WJ, Siefring GE, Jr. 1976. Streptokinase. *Methods Enzymol* 45:244–257. [https://doi.org/10.1016/s0076-6879\(76\)45024-3](https://doi.org/10.1016/s0076-6879(76)45024-3).
104. Castellino FJ, Sodetz JM. 1976. Rabbit plasminogen and plasmin isozymes. *Methods Enzymol* 45:273–286. [https://doi.org/10.1016/s0076-6879\(76\)45026-7](https://doi.org/10.1016/s0076-6879(76)45026-7).
105. Radek JT, Castellino FJ. 1989. Conformational properties of streptokinase. *J Biol Chem* 264:9915–9922. [https://doi.org/10.1016/S0021-9258\(18\)81747-1](https://doi.org/10.1016/S0021-9258(18)81747-1).
106. Cederholm-Williams SA, De Cock F, Lijnen HR, Collen D. 1979. Kinetics of the reactions between streptokinase, plasmin and alpha 2-antiplasmin. *Eur J Biochem* 100:125–132. <https://doi.org/10.1111/j.1432-1033.1979.tb02040.x>.
107. Lathem WW, Price PA, Miller VL, Goldman WE. 2007. A plasminogen-activating protease specifically controls the development of primary pneumonic plague. *Science* 315:509–513. <https://doi.org/10.1126/science.1137195>.
108. Claes H, Vermeylen J. 1974. Physico-chemical and proenzyme properties of NH₂-terminal glutamic acid and NH₂-terminal lysine human plasminogen. Influence of 6-aminohexanoic acid. *Biochim Biophys Acta* 342:351–359. [https://doi.org/10.1016/0005-2795\(74\)90090-7](https://doi.org/10.1016/0005-2795(74)90090-7).
109. Bokarewa MI, Jin T, Tarkowski A. 2006. *Staphylococcus aureus*: staphylokinase. *Int J Biochem Cell Biol* 38:504–509. <https://doi.org/10.1016/j.biocel.2005.07.005>.
110. Lahteenmaki K, Kuusela P, Korhonen TK. 2001. Bacterial plasminogen activators and receptors. *FEMS Microbiol Rev* 25:531–552. <https://doi.org/10.1111/j.1574-6976.2001.tb00590.x>.
111. Ullberg M, Kronvall G, Karlsson I, Wiman B. 1990. Receptors for human plasminogen on Gram-negative bacteria. *Infect Immun* 58:21–25. <https://doi.org/10.1128/IAI58.1.21-25.1990>.
112. Ullberg M, Kuusela P, Kristiansen BE, Kronvall G. 1992. Binding of plasminogen to *Neisseria meningitidis* and *Neisseria gonorrhoeae* and formation of surface-associated plasmin. *J Infect Dis* 166:1329–1334. <https://doi.org/10.1093/infdis/166.6.1329>.
113. Ringner M, Valkonen KH, Wadstrom T. 1994. Binding of vitronectin and plasminogen to *Helicobacter pylori*. *FEMS Immunol Med Microbiol* 9:29–34. <https://doi.org/10.1111/j.1574-695X.1994.tb00470.x>.
114. Khin MM, Ringner M, Aleljung P, Wadstrom T, Ho B. 1996. Binding of human plasminogen and lactoferrin by *Helicobacter pylori* coccoid forms. *J Med Microbiol* 45:433–439. <https://doi.org/10.1099/00222615-45-6-433>.
115. Parkkinen J, Korhonen TK. 1989. Binding of plasminogen to *Escherichia coli* adhesion proteins. *FEBS Lett* 250:437–440. [https://doi.org/10.1016/0014-5793\(89\)80772-0](https://doi.org/10.1016/0014-5793(89)80772-0).
116. Sjobring U, Pohl G, Olsen A. 1994. Plasminogen, absorbed by *Escherichia coli* expressing curli or by *Salmonella enteritidis* expressing thin aggregative fimbriae, can be activated by simultaneously captured tissue-type plasminogen activator (t-PA). *Mol Microbiol* 14:443–452. <https://doi.org/10.1111/j.1365-2958.1994.tb02179.x>.
117. Baranton G, Assous M, Postic D. 1992. Three bacterial species associated with Lyme borreliosis. Clinical and diagnostic implications. *Bull Acad Natl Med* 176:1075–1085. (In French.)
118. Grosskinsky S, Schott M, Brenner C, Cutler SJ, Kracicz P, Zipfel PF, Simon MM, Wallich R. 2009. *Borrelia recurrentis* employs a novel multifunctional surface protein with anti-complement, anti-opsonic and invasive potential to escape innate immunity. *PLoS One* 4:e4858. <https://doi.org/10.1371/journal.pone.0004858>.
119. Hovis KM, Freedman JC, Zhang H, Forbes JL, Marconi RT. 2008. Identification of an antiparallel coiled-coil/loop domain required for ligand binding by the *Borrelia hermsii* FhbA protein: additional evidence for the

- role of FhbA in the host-pathogen interaction. *Infect Immun* 76:2113–2122. <https://doi.org/10.1128/IAI.01266-07>.
120. Hammerschmidt C, Koenigs A, Siegel C, Hallstrom T, Skerka C, Wallich R, Zipfel PF, Kraiczy P. 2014. Versatile roles of CspA orthologs in complement inactivation of serum-resistant Lyme disease spirochetes. *Infect Immun* 82:380–392. <https://doi.org/10.1128/IAI.01094-13>.
 121. Barbour AG, Campeau Miller S. 2014. Genome sequence of *Borrelia parkeri*, an agent of enzootic relapsing fever in western North America. *Genome Announc* 2:e00018-14. <https://doi.org/10.1128/genomeA.00018-14>.
 122. Barbour AG. 2016. Chromosome and plasmids of the tick-borne relapsing fever agent *Borrelia hermsii*. *Genome Announc* 4:e00528-16. <https://doi.org/10.1128/genomeA.00528-16>.
 123. Welsh J, Pretzman C, Postic D, Saint Girons I, Baranton G, McClelland M. 1992. Genomic fingerprinting by arbitrarily primed polymerase chain reaction resolves *Borrelia burgdorferi* into three distinct phyletic groups. *Int J Syst Bacteriol* 42:370–377. <https://doi.org/10.1099/00207713-42-3-370>.
 124. Foldvari G, Farkas R, Lakos A. 2005. *Borrelia spielmanii* erythema migrans, Hungary. *Emerg Infect Dis* 11:1794–1795. <https://doi.org/10.3201/eid1111.050542>.
 125. Richter D, Postic D, Sertour N, Livey I, Matuschka FR, Baranton G. 2006. Delineation of *Borrelia burgdorferi* sensu lato species by multilocus sequence analysis and confirmation of the delineation of *Borrelia spielmanii* sp. nov. *Int J Syst Evol Microbiol* 56:873–881. <https://doi.org/10.1099/ijs.0.64050-0>.
 126. Seling A, Siegel C, Fingerle V, Jutras BL, Brissette CA, Skerka C, Wallich R, Zipfel PF, Stevenson B, Kraiczy P. 2010. Functional characterization of *Borrelia spielmanii* outer surface proteins that interact with distinct members of the human factor H protein family and with plasminogen. *Infect Immun* 78:39–48. <https://doi.org/10.1128/IAI.00691-09>.
 127. Platonov AE, Karan LS, Kolyasnikova NM, Makhneva NA, Toporkova MG, Maleev VV, Fish D, Krause PJ. 2011. Humans infected with relapsing fever spirochete *Borrelia miyamotoi*, Russia. *Emerg Infect Dis* 17:1816–1823. <https://doi.org/10.3201/eid1710.101474>.
 128. Telford SR, III, Molloy PJ, Berardi VP. 2015. *Borrelia miyamotoi*. *Ann Intern Med* 163:963–964. <https://doi.org/10.7326/L15-5187>.
 129. Jobe DA, Lovrich SD, Oldenburg DG, Kowalski TJ, Callister SM. 2016. *Borrelia miyamotoi* infection in patients from upper midwestern United States, 2014–2015. *Emerg Infect Dis* 22:1471–1473. <https://doi.org/10.3201/eid2208.151878>.
 130. Stone BL, Brissette CA. 2017. Host immune evasion by Lyme and relapsing fever borreliae: findings to lead future studies for *Borrelia miyamotoi*. *Front Immunol* 8:12. <https://doi.org/10.3389/fimmu.2017.00012>.
 131. Steere AC. 1989. Lyme disease. *N Engl J Med* 321:586–596. <https://doi.org/10.1056/NEJM19890813210906>.
 132. Garcia-Monco JC, Benach JL. 1995. Lyme neuroborreliosis. *Ann Neurol* 37:691–702. <https://doi.org/10.1002/ana.410370602>.
 133. Garcia-Monco JC, Benach JL. 2019. Lyme neuroborreliosis: clinical outcomes, controversy, pathogenesis, and polymicrobial infections. *Ann Neurol* 85:21–31. <https://doi.org/10.1002/ana.25389>.
 134. Szczepanski A, Furie MB, Benach JL, Lane BP, Fleit HB. 1990. Interaction between *Borrelia burgdorferi* and endothelium *in vitro*. *J Clin Invest* 85:1637–1647. <https://doi.org/10.1172/JCI114615>.
 135. Moriarty TJ, Shi M, Lin YP, Ebady R, Zhou H, Odisho T, Hardy PO, Salman-Dilgimen A, Wu J, Weening EH, Skare JT, Kubes P, Leong J, Chaconas G. 2012. Vascular binding of a pathogen under shear force through mechanistically distinct sequential interactions with host macromolecules. *Mol Microbiol* 86:1116–1131. <https://doi.org/10.1111/mmi.12045>.
 136. Coleman JL, Benach JL. 1999. Use of the plasminogen activation system by microorganisms. *J Lab Clin Med* 134:567–576. [https://doi.org/10.1016/s0022-2143\(99\)90095-1](https://doi.org/10.1016/s0022-2143(99)90095-1).
 137. Zimmer G, Schaible UE, Kramer MD, Mall G, Museteanu C, Simon MM. 1990. Lyme carditis in immunodeficient mice during experimental infection of *Borrelia burgdorferi*. *Virchows Arch A Pathol Anat Histopathol* 417:129–135. <https://doi.org/10.1007/BF02190530>.
 138. Barthold SW, Persing DH, Armstrong AL, Peeples RA. 1991. Kinetics of *Borrelia burgdorferi* dissemination and evolution of disease after intradermal inoculation of mice. *Am J Pathol* 139:263–273.
 139. Defosse DL, Duray PH, Johnson RC. 1992. The NIH-3 immunodeficient mouse is a model for Lyme borreliosis myositis and carditis. *Am J Pathol* 141:3–10.
 140. Barthold SW, de Souza MS, Janotka JL, Smith AL, Persing DH. 1993. Chronic Lyme borreliosis in the laboratory mouse. *Am J Pathol* 143:959–971.
 141. Ruderman EM, Kerr JS, Telford SR, III, Spielman A, Glimcher LH, Gravalles EM. 1995. Early murine Lyme carditis has a macrophage predominance and is independent of major histocompatibility complex class II-CD4+ T cell interactions. *J Infect Dis* 171:362–370. <https://doi.org/10.1093/infdis/171.2.362>.
 142. Coleman JL, Sellati TJ, Testa JE, Kew RR, Furie MB, Benach JL. 1995. *Borrelia burgdorferi* binds plasminogen, resulting in enhanced penetration of endothelial monolayers. *Infect Immun* 63:2478–2484. <https://doi.org/10.1128/IAI.63.7.2478-2484.1995>.
 143. Gebbia JA, Monco JC, Degen JL, Bugge TH, Benach JL. 1999. The plasminogen activation system enhances brain and heart invasion in murine relapsing fever borreliosis. *J Clin Invest* 103:81–87. <https://doi.org/10.1172/JCI5171>.
 144. Klempner MS, Noring R, Epstein MP, McCloud B, Rogers RA. 1996. Binding of human urokinase type plasminogen activator and plasminogen to *Borrelia* species. *J Infect Dis* 174:97–104. <https://doi.org/10.1093/infdis/174.1.97>.
 145. Nguyen NTT, Rottgerding F, Devraj G, Lin YP, Koenigs A, Kraiczy P. 2018. The complement binding and inhibitory protein CbiA of *Borrelia miyamotoi* degrades extracellular matrix components by interacting with plasmin(ogen). *Front Cell Infect Microbiol* 8:23. <https://doi.org/10.3389/fcimb.2018.00023>.
 146. Anda P, Gebbia JA, Backenson PB, Coleman JL, Benach JL. 1996. A glyceraldehyde-3-phosphate dehydrogenase homolog in *Borrelia burgdorferi* and *Borrelia hermsii*. *Infect Immun* 64:262–268. <https://doi.org/10.1128/IAI.64.1.262-268.1996>.
 147. Nordstrand A, Shamaei-Tousi A, Ny A, Bergstrom S. 2001. Delayed invasion of the kidney and brain by *Borrelia crocidurae* in plasminogen-deficient mice. *Infect Immun* 69:5832–5839. <https://doi.org/10.1128/iai.69.9.5832-5839.2001>.
 148. Klempner MS, Noring R, Epstein MP, McCloud B, Hu R, Limentani SA, Rogers RA. 1995. Binding of human plasminogen and urokinase-type plasminogen activator to the Lyme disease spirochete, *Borrelia burgdorferi*. *J Infect Dis* 171:1258–1265. <https://doi.org/10.1093/infdis/171.5.1258>.
 149. Fuchs H, Wallich R, Simon MM, Kramer MD. 1994. The outer surface protein A of the spirochete *Borrelia burgdorferi* is a plasmin(ogen) receptor. *Proc Natl Acad Sci U S A* 91:12594–12598. <https://doi.org/10.1073/pnas.91.26.12594>.
 150. Koenigs A, Hammerschmidt C, Jutras BL, Pogoryelov D, Barthel D, Skerka C, Kugelstadt D, Wallich R, Stevenson B, Zipfel PF, Kraiczy P. 2013. BBA70 of *Borrelia burgdorferi* is a novel plasminogen-binding protein. *J Biol Chem* 288:25229–25243. <https://doi.org/10.1074/jbc.M112.413872>.
 151. Coleman JL, Roemer EJ, Benach JL. 1999. Plasmin-coated *Borrelia burgdorferi* degrades soluble and insoluble components of the mammalian extracellular matrix. *Infect Immun* 67:3929–3936. <https://doi.org/10.1128/IAI.67.8.3929-3936.1999>.
 152. Hu LT, Perides G, Noring R, Klempner MS. 1995. Binding of human plasminogen to *Borrelia burgdorferi*. *Infect Immun* 63:3491–3496. <https://doi.org/10.1128/IAI.63.9.3491-3496.1995>.
 153. Caimano MJ, Eggers CH, Gonzalez CA, Radolf JD. 2005. Alternate sigma factor RpoS is required for the *in vivo*-specific repression of *Borrelia burgdorferi* plasmid lp54-borne ospA and lp6.6 genes. *J Bacteriol* 187:7845–7852. <https://doi.org/10.1128/JB.187.22.7845-7852.2005>.
 154. Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA. 1995. Induction of an outer surface protein on *Borrelia burgdorferi* during tick feeding. *Proc Natl Acad Sci U S A* 92:2909–2913. <https://doi.org/10.1073/pnas.92.7.2909>.
 155. Coleman JL, Gebbia JA, Piesman J, Degen JL, Bugge TH, Benach JL. 1997. Plasminogen is required for efficient dissemination of *B. burgdorferi* in ticks and for enhancement of spirochetemia in mice. *Cell* 89:1111–1119. [https://doi.org/10.1016/s0092-8674\(00\)80298-6](https://doi.org/10.1016/s0092-8674(00)80298-6).
 156. Xu Q, McShan K, Liang FT. 2008. Essential protective role attributed to the surface lipoproteins of *Borrelia burgdorferi* against innate defences. *Mol Microbiol* 69:15–29. <https://doi.org/10.1111/j.1365-2958.2008.06264.x>.
 157. Lagal V, Portnoi D, Faure G, Postic D, Baranton G. 2006. *Borrelia burgdorferi* sensu stricto invasiveness is correlated with OspC-plasminogen affinity. *Microbes Infect* 8:645–652. <https://doi.org/10.1016/j.micinf.2005.08.017>.
 158. Onder O, Humphrey PT, McOmber B, Korobova F, Francella N, Greenbaum DC, Brisson D. 2012. OspC is potent plasminogen receptor on surface of *Borrelia burgdorferi*. *J Biol Chem* 287:16860–16868. <https://doi.org/10.1074/jbc.M111.290775>.
 159. Brissette CA, Haupt K, Barthel D, Cooley AE, Bowman A, Skerka C, Wallich R, Zipfel PF, Kraiczy P, Stevenson B. 2009. *Borrelia burgdorferi*

- infection-associated surface proteins ErpP, ErpA, and ErpC bind human plasminogen. *Infect Immun* 77:300–306. <https://doi.org/10.1128/IAI.01133-08>.
160. Krawczyk P, Skerka C, Kirschfink M, Brade V, Zipfel PF. 2001. Immune evasion of *Borrelia burgdorferi* by acquisition of human complement regulators FHL-1/reconectin and Factor H. *Eur J Immunol* 31:1674–1684. [https://doi.org/10.1002/1521-4141\(200106\)31:6<1674::AID-IMMU1674>3.0.CO;2-2](https://doi.org/10.1002/1521-4141(200106)31:6<1674::AID-IMMU1674>3.0.CO;2-2).
 161. Hallstrom T, Haupt K, Krawczyk P, Hortschansky P, Wallich R, Skerka C, Zipfel PF. 2010. Complement regulator-acquiring surface protein 1 of *Borrelia burgdorferi* binds to human bone morphogenetic protein 2, several extracellular matrix proteins, and plasminogen. *J Infect Dis* 202:490–498. <https://doi.org/10.1086/653825>.
 162. Brissette CA, Cooley AE, Burns LH, Riley SP, Verma A, Woodman ME, Bykowski T, Stevenson B. 2008. Lyme borreliosis spirochete Erp proteins, their known host ligands, and potential roles in mammalian infection. *Int J Med Microbiol* 298 Suppl 1:257–267. <https://doi.org/10.1016/j.ijmm.2007.09.004>.
 163. Rossmann E, Krawczyk P, Herzberger P, Skerka C, Kirschfink M, Simon MM, Zipfel PF, Wallich R. 2007. Dual binding specificity of a *Borrelia hermsii*-associated complement regulator-acquiring surface protein for factor H and plasminogen discloses a putative virulence factor of relapsing fever spirochetes. *J Immunol* 178:7292–7301. <https://doi.org/10.4049/jimmunol.178.11.7292>.
 164. Schott M, Grosskinsky S, Brenner C, Krawczyk P, Wallich R. 2010. Molecular characterization of the interaction of *Borrelia parkeri* and *Borrelia turicatae* with human complement regulators. *Infect Immun* 78:2199–2208. <https://doi.org/10.1128/IAI.00089-10>.
 165. Hovis KM, McDowell JV, Griffin L, Marconi RT. 2004. Identification and characterization of a linear-plasmid-encoded factor H-binding protein (FhbA) of the relapsing fever spirochete *Borrelia hermsii*. *J Bacteriol* 186:2612–2618. <https://doi.org/10.1128/jb.186.9.2612-2618.2004>.
 166. Redlitz A, Fowler BJ, Plow EF, Miles LA. 1995. The role of an enolase-related molecule in plasminogen binding to cells. *Eur J Biochem* 227:407–415. <https://doi.org/10.1111/j.1432-1033.1995.tb20403.x>.
 167. Lottenberg R, Broder CC, Boyle MD. 1987. Identification of a specific receptor for plasmin on a group A streptococcus. *Infect Immun* 55:1914–1918. <https://doi.org/10.1128/IAI.55.8.1914-1918.1987>.
 168. Miles LA, Dahlberg CM, Plescia J, Felez J, Kato K, Plow EF. 1991. Role of cell-surface lysines in plasminogen binding to cells: identification of alpha-enolase as a candidate plasminogen receptor. *Biochemistry* 30:1682–1691. <https://doi.org/10.1021/bi00220a034>.
 169. Pancholi V. 2001. Multifunctional alpha-enolase: its role in diseases. *Cell Mol Life Sci* 58:902–920. <https://doi.org/10.1007/pl00000910>.
 170. Floden AM, Watt JA, Brissette CA. 2011. *Borrelia burgdorferi* enolase is a surface-exposed plasminogen binding protein. *PLoS One* 6:e27502. <https://doi.org/10.1371/journal.pone.0027502>.
 171. Toledo A, Coleman JL, Kuhlow CJ, Crowley JT, Benach JL. 2012. The enolase of *Borrelia burgdorferi* is a plasminogen receptor released in outer membrane vesicles. *Infect Immun* 80:359–368. <https://doi.org/10.1128/IAI.05836-11>.
 172. Nogueira SV, Smith AA, Qin JH, Pal U. 2012. A surface enolase participates in *Borrelia burgdorferi*-plasminogen interaction and contributes to pathogen survival within feeding ticks. *Infect Immun* 80:82–90. <https://doi.org/10.1128/IAI.05671-11>.
 173. Nowalk AJ, Nolder C, Clifton DR, Carroll JA. 2006. Comparative proteome analysis of subcellular fractions from *Borrelia burgdorferi* by NEPHGE and IPG. *Proteomics* 6:2121–2134. <https://doi.org/10.1002/pmic.200500187>.
 174. Fulde M, Rohde M, Polok A, Preissner KT, Chhatwal GS, Bergmann S. 2013. Cooperative plasminogen recruitment to the surface of *Streptococcus canis* via M protein and enolase enhances bacterial survival. *mBio* 4:e00629-12. <https://doi.org/10.1128/mBio.00629-12>.
 175. Hu LT, Pratt SD, Perides G, Katz L, Rogers RA, Klempner MS. 1997. Isolation, cloning, and expression of a 70-kilodalton plasminogen binding protein of *Borrelia burgdorferi*. *Infect Immun* 65:4989–4995. <https://doi.org/10.1128/IAI.65.12.4989-4995.1997>.
 176. Gebbia JA, Coleman JL, Benach JL. 2001. *Borrelia burgdorferi* upregulate release and activation of matrix metalloproteinase gelatinase B (MMP-9) and collagenase 1 (MMP-1) in human cells. *Infect Immun* 69:456–462. <https://doi.org/10.1128/IAI.69.1.456-462.2001>.
 177. Gebbia JA, Coleman JL, Benach JL. 2004. Selective induction of matrix metalloproteinases by *Borrelia burgdorferi* via toll-like receptor 2 in monocytes. *J Infect Dis* 189:113–119. <https://doi.org/10.1086/380414>.
 178. Behera AK, Thorpe CM, Kidder JM, Smith W, Hildebrand E, Hu LT. 2004. *Borrelia burgdorferi*-induced expression of matrix metalloproteinases from human chondrocytes requires mitogen-activated protein kinase and Janus kinase/signal transducer and activator of transcription signaling pathways. *Infect Immun* 72:2864–2871. <https://doi.org/10.1128/iai.72.5.2864-2871.2004>.
 179. Behera AK, Hildebrand E, Scagliotti J, Steere AC, Hu LT. 2005. Induction of host matrix metalloproteinases by *Borrelia burgdorferi* differs in human and murine Lyme arthritis. *Infect Immun* 73:126–134. <https://doi.org/10.1128/IAI.73.1.126-134.2005>.
 180. Behera AK, Hildebrand E, Uematsu S, Akira S, Coburn J, Hu LT. 2006. Identification of a TLR-independent pathway for *Borrelia burgdorferi*-induced expression of matrix metalloproteinases and inflammatory mediators through binding to integrin alpha 3 beta 1. *J Immunol* 177:657–664. <https://doi.org/10.4049/jimmunol.177.1.657>.
 181. Perides G, Tanner-Brown LM, Eskildsen MA, Klempner MS. 1999. *Borrelia burgdorferi* induces matrix metalloproteinases by neural cultures. *J Neurosci Res* 58:779–790. [https://doi.org/10.1002/\(SICI\)1097-4547\(19991215\)58:6<779::AID-JNRS>3.0.CO;2-L](https://doi.org/10.1002/(SICI)1097-4547(19991215)58:6<779::AID-JNRS>3.0.CO;2-L).
 182. Lin B, Kidder JM, Noring R, Steere AC, Klempner MS, Hu LT. 2001. Differences in synovial fluid levels of matrix metalloproteinases suggest separate mechanisms of pathogenesis in Lyme arthritis before and after antibiotic treatment. *J Infect Dis* 184:174–180. <https://doi.org/10.1086/322000>.
 183. Zhao Z, Chang H, Trevino RP, Whren K, Bhawan J, Klempner MS. 2003. Selective up-regulation of matrix metalloproteinase-9 expression in human erythema migrans skin lesions of acute Lyme disease. *J Infect Dis* 188:1098–1104. <https://doi.org/10.1086/379039>.
 184. Zhao Z, Fleming R, McCloud B, Klempner MS. 2007. CD14 mediates cross talk between mononuclear cells and fibroblasts for upregulation of matrix metalloproteinase 9 by *Borrelia burgdorferi*. *Infect Immun* 75:3062–3069. <https://doi.org/10.1128/IAI.00202-07>.
 185. Coleman JL, Gebbia JA, Benach JL. 2001. *Borrelia burgdorferi* and other bacterial products induce expression and release of the urokinase receptor (CD87). *J Immunol* 166:473–480. <https://doi.org/10.4049/jimmunol.166.1.473>.
 186. Coleman JL, Benach JL. 2003. The urokinase receptor can be induced by *Borrelia burgdorferi* through receptors of the innate immune system. *Infect Immun* 71:5556–5564. <https://doi.org/10.1128/iai.71.10.5556-5564.2003>.
 187. Hovius JW, Bijlsma MF, van der Windt GJ, Wiersinga WJ, Boukens BJ, Coumou J, Oei A, de Beer R, de Vos AF, van 't Veer C, van Dam AP, Wang P, Fikrig E, Levi MM, Roelofs JJ, van der Poll T. 2009. The urokinase receptor (uPAR) facilitates clearance of *Borrelia burgdorferi*. *PLoS Pathog* 5:e1000447. <https://doi.org/10.1371/journal.ppat.1000447>.
 188. Flick MJ, Bugge TH. 2017. Plasminogen-receptor KT: plasminogen activation and beyond. *J Thromb Haemost* 15:150–154. <https://doi.org/10.1111/jth.13541>.
 189. Andronikos NM, Chen EI, Baik N, Bai H, Parmer CM, Kiosses WB, Kamps MP, Yates JR, III, Parmer RJ, Miles LA. 2010. Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-RKT, a major regulator of cell surface plasminogen activation. *Blood* 115:1319–1330. <https://doi.org/10.1182/blood-2008-11-188938>.
 190. Lighvani S, Baik N, Diggs JE, Khaldoyanidi S, Parmer RJ, Miles LA. 2011. Regulation of macrophage migration by a novel plasminogen receptor Plg-R KT. *Blood* 118:5622–5630. <https://doi.org/10.1182/blood-2011-03-344242>.
 191. Thaler B, Baik N, Hohensinner PJ, Baumgartner J, Panzenbock A, Stojkovic S, Demyanets S, Huk I, Rega-Kaun G, Kaun C, Prager M, Fischer MB, Huber K, Speidl WS, Parmer RJ, Miles LA, Wojta J. 2019. Differential expression of Plg-RKT and its effects on migration of proinflammatory monocyte and macrophage subsets. *Blood* 134:561–567. <https://doi.org/10.1182/blood.2018850420>.
 192. Hallstrom T, Siegel C, Morgelin M, Krawczyk P, Skerka C, Zipfel PF. 2013. CspA from *Borrelia burgdorferi* inhibits the terminal complement pathway. *mBio* 4:e00481-13. <https://doi.org/10.1128/mBio.00481-13>.